

# INDEX\*.

- Aconiopteris obtusa*, 578.
- Acrophorus assamicus*, 443; *Hookeri*, 443; *immersus*, 443; *nodosus*, 444; *pseudo-cystopteris*, 444; *pulcher*, 444; *Thomsoni*, 443.
- Acropteris australis*, 505; *radiata*, 505.
- Acrospira* (char. gen.), 255; *asphodeloides*, 255.
- Acrostichum alaicorne*, 595; *angulatum*, 576; *appendiculatum*, 577, 578, 596, 603, 604, var. *Hamiltoniana*, 578; *asplenifolium*, 577; *aureum*, 582, 596; *australe*, 505; *axillare*, 579, 596, 601; *Blumeanum*, 580; *conforme*, 576, 577, 595; *contaminans*, 596; *contractum*, 603; *coriaceum*, 595; *costatum*, 580, 581, (*Wall.*) 596, var. (*Hk.*) 572; *crispatum*, 580, 581, 596, var. *Blumeana*, 580, var. *contaminans*, 580, 596, var. *pseud-undulata*, 581; *danæfolium*, 582; *decurrens*, 595; *dichotomum*, 505; *diversifolium*, 580; *emarginatum*, 582; *Finlaysonianum*, 603; *flagelliferum*, 579, 581, 596; *fuciforme*, 595; *Gayanum*, 576; *glandulosum*, 576; *gorgoneum*, (*Kaulf.*) 578, (*Bl.*) 576; *Hamiltonianum*, 577, 596; *heterophyllum*, 575; *hybridum*, 595; *inæquale*, 582; *laurifolium*, 576; *ludens*, 577, 578, 605; *Marantæ*, 567; *marginatum*, 576, 595; *minus*, 579; *neriifolium*, 577, 595; *numulariæfolium*, 554; *obovatum*, 554; *palustre*, 577; *proliferum*, 581; *punctatum*, 561; *punctulatum*, 596; *radiatum*, 505; *repandum*, 580; *rigidum*, 596; *rivulare*, 579, 603; *scandens*, 577, 596; *septentrionale*, 478; *setosum*, 577, 596; *speciosum*, 582; *spicatum*, 582; *stelligerum*, 577, 604; *stigmatolepis*, 577; *subcrenatum*, 581; *terminans*, 581, 604; *thalictroides*, 471; *Thelypteris*, 517; *tricuspe*, 581; *triquetrum*, 473, 596; *variabile*, 578, 579, var. *lanceolata*, 579; *vestitum*, 603; *virens*, (*Wall.*) 581, 598, 603, 604, (*Bk.*) 580, (*Hk. & Grev.*) 581, var. (*Hk.*) 580, var. (*Hk. & Bk.*) 572, *viscosum*, 577, 581, 595; *viviparum*, 577, 578, 596; *Wightianum*, 577, 603.
- Actiniopteris dichotoma*, 505; *radiata*, 505.
- Actinostachys digitata*, 583, 595.
- Adenotheca aphylla* (*Welw.*), 255.
- Adiantum amœnum*, 454, 597; *capillus*, 453; *Capillus-Veneris*, 453, 597; *caudatum*, 453, 597, var. *rhizophorum*, 453; *ciliatum*, 453; *dolabriforme*, 452; *Edgworthii*, 453; *filiforme*, 452; *flabellatum*, 454, 597, 604; *flagelliferum*, 453, 597; *hirsutum*, 453, 604; *hispidulum*, 454, 604; *lunulatum*, 452, 597; *microphyllum*, 453; *pedatum*, 453; *proliferum*, 453; *reni-*
- forme*, 597; *repens*, 442; *rhizophorum*, 597; *scabrum*, 597; *soboliferum*, 453, 597; *tenerum*, 453; *venustum*, 453, 597; *vestitum*, 453.
- Adoxa*, structure of, Rev. G. Henslow on, 177.
- Æstivations, origin of floral, 177.
- Agaricini*, 119; of Brisbane, 399.
- Agaricus fimicola*, 119; *Gardneri*, 399; *junceus*, 119; *semiorbicularis*, 119; *spadicens*, 119; *sylvaticus*, 119.
- Agasta*, 59; *asiatica*, 61; *indica*, 63; *splendida*, 60.
- Aglaia odorata*, distrib. of, 233.
- Abuta* = *Agasta splendida*, 60.
- Aira*, awn of, 164.
- Albinism in Balsams, Dr. Lowne's exper. on, 328.
- Albica angolensis*, 251; *chlorantha*, 251; *galeata*, 251; *monophylla*, 251; *myogaloides*, 250; *subspicata*, 251.
- Alismacæ*, self-fertilization of, 378, 393.
- Allantodia denticulata*, 490; *incisa*, 490; *javanica*, 505, 599; *sylvatica*, 501; *tenella*, 491.
- Allium angolense*, 262.
- Allosorus calomelanos*, 461; *crispus*, 459, 460; *gracilis*, 460; *Stolleri*, 460.
- Aloe andongensis*, 263; *angolensis*, 263; *littoralis*, 263; *palmiformis*, 263; *platyphylla*, 264; *zebrina*, 264.

\* The names of species adopted by the author precede synonyms; the latter are in alphabetical order.

- Alsophila Andersoni*, 433; *Brunoniana*, 427, 430, 431, 605, (*Bedd.*) 432; *comosa*, 433; *contaminans*, 432; *decipiens*, 430; *gigantea*, 433; *glabra*, 433, 496; *glauca*, 431, 433, 605; *glaucescens*, 432, 605; *Grevilleana*, 449, 605; *Helforiana*, 433; *hasiana*, 432; *latebrosa*, 429, (*Bedd.*) 432, (*Ilk.*) 431, var. *hemitelioides* (*J. Scott*) 430, var. *Schmidiana* (*Kze.*) 432; *Metteniana*, 433; *Oldhami*, 432, 433; *ornata*, 432; *Scottiana*, 433; *Sollyana*, 432; *Wallichiana*, 432.
- Ampelopteris elegans*, 548; *firma*, 548.
- Amphicosmia decipiens*, 430.
- Androscepiæ arundinacea*, awn of, 164.
- Anemia flexuosa*, 596.
- Anemone montana*, burying mechanism of achene of, 164; torsion of awn of, 158.
- Angiopteris crassipes*, 585, 599; *evecta*, 585, 599.
- Angolan Herbarium of Welwitsch, J. G. Baker on the Liliaceæ, Iridaceæ, Hypoxidaceæ, and Hamodoraceæ of, 245.
- Anisogonium* (subgen.), 503; *esculentum*, 503; *heterophlebium*, 503; *serampurense*, 503; *sylvaticum*, 498.
- Anisostemon* (sect. *Bl.*), 113.
- Anthericum andongense*, 257; *arenarium*, 259; *benguellense*, 257; *calyptrocarpum*, 258; *colubrinum*, (*Welw.*) 256; *dissitiflorum*, 257; *drimiopsis*, 259; *falcatum*, (*Welw.*) 256; *Jacquinianum*, 259, var. *affinis*, 259; *limosum*, 257; *molle*, 259; *orchideum*, 258; *pleiostachyum*, (*Welw.*) 255; *pteroaulon*, 258; *pyrenicarpum*, 259; *tenellum*, 256; *ustulatum*, 258.
- Anthesteria ciliata*, awn of, 164.
- Antholyza huillensis*, 270.
- Antrophyum Boryanum*, 573, 596; *coriaceum*, (*Wall.*) 572, 596, (*Bl.*) 570; *falcatum*, 573; *involutum*, 570; *lanceolatum*, 570; *latifolium*, 572, 573; *parvulum*, 573; *plantagineum*, 572, 573; *plicatum*, 572; *pumilum*, 596; *reticulatum*, (*Kaulf.*) 573, 596, (*Bedd.*) 572, (*Fée*) 572, (*Wall.*) 572, var. *parvula*, 573.
- Apocynaceæ, plumes of seed of, 165.
- Appunia* (*Hook. f.*), 169.
- Areyria umbrina*, 128.
- Aristea angolensis*, 270.
- Arrhenatherum*, awn of, 164.
- Arthobotrys Avana*, 522, 603; *macrocarpa*, 522, 603.
- Arthronia punctilliformis*, 146.
- Asparagus africanus*, 254; *angolensis*, 254; *benguellensis*, 253; *defloxus*, 254; *drepanophyllus*, 254; *equisetoides*, 253; *psilurus*, 253; *pubescens*, 254; *racemosus*, 254.
- Aspergillus glaucus*, 406.
- Aspidium abruptum*, 534; *aculeatum*, 508, 509, (*Swtz.*) var. 603, var. *biaristata*, 510, var. *biaristatum* f. *Khasiana*, 511, var. *lobata*, 509, var. *mucronifolia*, 509, var. *rufobarbata*, 509, var. *semifertilis*, 509, var. *setosa*, 510; *acutum*, 541; *adnatum*, 518; *aduncum*, 603; *affine*, 509, 603; *alatum*, 536, 603; *amabile*, 510, 602; *amboinense*, 532; *angustifrons*, 514; *anomophyllum*, 512; *apiciflorum*, 518, 602; *appendiculatum*, 516, 543, 602; *aridum*, 531; *aristatum*, 511, 529, 602, var. *affinis*, 511, var. *assamica*, 511; *articulatum*, 603; *Athyrium*, 491; *Atkinsoni*, 506; *atratum*, 513, 603; *auriculatum*, (*Swtz.*) 506, 602, 603, 605, (*Don*) 507, (*Wall.*) 540, var. *cæspitosa*, 507, var. *lenta*, 507, var. *marginata*, 507, var. *normalis*, 507; *biaristatum*, 511; *Brunonianum*, 522, 602; *caducum*, 512, 603, 605; *cataphoron*, 524; *cæspitosum*, 603; *canariense*, 521; *canescens*, 533, 602; *canum*, 515, 533, 603; *caruifolium*, 511; *caryotideum*, 512, 603; *cicutarium*, 539; *ciliatum*, 515, 602; *coadunatum*, 540, 603; *coniifolium*, 511, 602; *conioneuron*, 529; *conjugatum*, 535; *cordifolium*, 540; *coriaceum*, 605; *cornu-cervi*, 511; *crinitum*, 602; *cristatum*, 519; *cucullatum*, 530; *cuspidatum*, 513; *decurrens*, 539; *densum*, 523, 603; *discretum*, 510; *divisum*, 527, 546, 603; *Donianum*, 520; *dropanopterum*, 493; *eburneum*, 486, 493, 523, 603; *elongatum*, 520; *eriocarpum*, 525, 602; *eusorum*, 534; *exaltatum*, 541, 603; *extensum*, 529; *falcatum*, 512, 603; *Filix-fœmina*, 491; *Filix-mas*, 519; *fimbriatum*, 494, 602; *Finlaysonianum*, 605; *flaccidum*, 513; *flagelliferum*, 605; *fœniculaceum*, 427, 434; *foliosum*, 434, 444, 602; *fontanum*, 484; *fuscipes*, 534, 602; *glanduliferum*, 516, 602; *glanduligerum*, 513; *gracilescens*, 513; *grandifolium*, 537; *Hamiltonianum*, 605; *hirsutulum*, (*Wall.*) 515, 605, (*Ham.*) 605, (*Schk.*) 541; *hirtipes*, 513; *ilicifolium*, 508, 603; *imbricatum*, 540; *intermedium*, 526; *Lachenense*, 506, 510; *Leuzeanum*, 535; *Lonchitis*, 505; *macrocarpum*, 489; *macrophyllum*, 539; *marginatum*, 496, 521, 602, 603; *Maximowiczianum*, 511; *megaphyllum*, 532; *membranifolium*, 535; *microcarpon*, 561; *molle*, 533; *multicaudatum*, 540; *multidentatum*, 443, 602; *multijugum*, 529, 533, 602; *multilineatum*, 534, 602; *musæfolium*, 542; *neriiforme*, 542; *nitidulum*, 603; *nodosum*, 444; *nymphale*, 530; *obliquum*, 507; *obtusifolium*, 540; *ocellatum*, 507, 602; *oethodes*, 516; *odoratum*, 525; *paleaceum*, 520; *palmipes*, 511; *paludosum*, 544; *parasiticum*, (*Herb. Madr.*) 605, (*Swtz.*) 533, 605; *patentissimum*, 520, 602; *pennigerum*, 532; *pilosulum*, 448, 525, 602; *pilosum*, 541; *polyblepharon*, 510; *polymorphum*, (*Bedd.*) 538, 603, (*Wall.*) 537, *Prescottianum*, 508, 510, 602, var. *Bakeriana*, 510, var. *castanea*, 510; *prionophyllum*, 529, 534, 602; *procurrens*, 530; *prolixum*, 516; *pteropus*, (*Kze.*) 539, (*Mett.*) 539; *puberum*, 448, 515, 602; *pungens*, 508, 603; *purpurascens*, 523, 524; *remotum*, 523; *repandum*, 537; *rhomboideum*, 602; *rigidum*, (*Swtz.*) 523, var. *australe* (*Mett.*) 521; *riparium*, 602; *rostratum*, 537, 603

- rufo-barbatum, 603; Schkuhrii, 541; semibipinnatum, 603; setosum, 603; siifolium, 538, 539; sikkimense, (*Bk.*) 427, (*Hk. & Bk.*) 525; Simonsii, 537; singaporianum, 603; solutum, 533, 602; sophoroides, 605; speciosum, 511; spectabile, (*Bl.*) 513, (*Wall.*) 435, 526, 603, splendens, (*Ham.*) 605, (*Willd.*) 541; squarrosus, (*Don*) 509, (*Wall.*) 489, 602; stimulans, 508; subconfluens, 536; subdiaphanum, 525, 602; subdigitatum, 546; sublanosum, 541, 602; syrmaticum, 518; Tavoyanum, 540, 603; tectum, 533, 603; Teijsmannianum, 538, 539; Telfairianum, 603; tenella, 491; terminans, 603; Thelypteris, 514, 517; Thomsoni, 508, 510, var. gracilis, 508; trifoliatum, 605; truncatum, 534; uliginosum, 520, 528; unitum, 529, 602; variolosum, 538, 603; vastum, 536; venulosum, 531, 602; vestitum, 510; Wallichianum, (*Spreng.*) 520, (*Wall.*) 542; Wallichii, 542; Weigleanum, 523; xylodes, 516; Zollingerianum, 538.
- Asplenium* abscissum, 481; acrostichoides, 488; acuminatum, 498, 500, 599; Adiantum-nigrum, 483, 485; alternans, 476, 600; alternifolium, 497; ambiguum, 498; amœnum, 481; anceps, 477; anthriscifolium, 482; arborescens, 503; argutans, 498; arifolium, 576; asperum, 502; aspidioides, 494; Atkinsoni, 487, var. Andersoni, 487; auritum, 600; australasicum, 475; australe, 505; bantamense, 497, 504, 599, 606; Beddomei, 479; bellum, 496, 502, 603; bipinnatum, 503; brevisorum, 494, 599; Brunoniana, 505; Brunonianum, 505; bulbiferum, 485, 599; bullatum, 485, 599; cæspitosum, 482, 599; castaneum, 477; caudatum, 600; Ceterach, 504; cheilosorum, 481; Clarkei, 489, 599; concinnum, 485, 599; contaminans, 604; contiguum, 479; coriaceum, (*Fée*) 476, (*Roxb.*) 480; costale, 490; crinicaule, 479, 480; cristatum, 481, 599; cultratum, 604; cuneatum, var. splendens, 484; Dalhousiæ, 476; decipiens, 489, 495; decurrens, (*Bedd.*) 503, (*Wall.*) 481, 599; decussatum, 499, 604; densum, 477; dentigerum, 492; depauperatum, 482, 600; dilatatum, 502; divaricatum, 604; diversifolium, 502, 599; Donianum, 497; drepanophyllum, 487; eburneum, 493; elatum, 498; elegans, 497; emarginato-dentatum, 481; ensiforme, 476, 595, 599, var. australis, 476; erectum, 478, var. trapeziforme, 479; eroso-dentatum, 481; erythrocaulon, 481; esculentum, 503, 599, 604; exiguum, 484; falcatum, 479, 480, (*Lamk.*) var. 599, 600, 603, var. caudatum, 480, var. macrophyllum, 480; fallax, 489; Filix-foemina, 489, 491, 600, 602, var. attenuata, 492, var. dentigera, 491, var. flabellulata, 493, var. Parasnathensis, 493, var. pectinata, 492, 495, var. polyspora, 493, var. retusa, 491, 492, var. tenella, 492; fimbriatum, (*Hk.*) 494, (*Kze.*) 485, 489, 491, (*Hk.*) var. 602, var. foliosa, 495, var. leptophyllum, 485, var. sphæropteroides, 495; Finlaysonianum, (*Hk.*), 504, 605, (*Wall.*) 480, 599; flagelliferum, 478, 599; foliolosum, 492, 604; fontanum, 484; fraxinifolium, 497, 599; frondosum, 503; furcatum, 484, 604; Gardneri, 480; Goringianum, 489; grammitoides, 499; Grevilleanum, 603; Grevillei, 476; Griffithianum, 476, var. australis, 476; Griffithii, 502; gymnogrammoides, (*Bedd.*) 490, (*Klotzsch*) 490; Halleri, 484; Hancei, 479; heterocarpum, 480, 481, 599; heterophlebium, 503; hirsutum, 484, 599; Hookerianum, 497, 504, 605, 606; humile, 483; insigne, 483; integerrimum, 504; japonicum, 498, 502, 604, var. chattagramica, 499, var.? Schkuhrii, 499; javanicum, 505; Jerdoni, 502; laciniatum, 481, 482, 599, 610, var. depauperata, 482, var. planicaule, 482; lætum, 481, 599; lanceolatum, 485; lan-
- ceum, 496, 599; laserpitiifolium, 484; lasiopteris, 499; latifolium, 498, 500, 502, 599, 600, var. decurrens, 503, var. dilatatum, 503, var. frondosum, 503, var. japonicum, 503; Lobbianum, 497; lobulosum, 478, 599; longifolium, 427, 478, 479, 599; longissimum, 478, 599; macrocarpum, 486, 488, 602, 603, var. Atkinsoni, 489, var. pinnata, 489, var. squarrosus, 493; macrophyllum, 480, 504; marginatum, 500, 604; maximum, (*Don*) 503, (*Hk. & Bk.*) 502; minus, 477; monanthemum, 477; Moritzii, 503; multicaudatum, 496, 502, 600, var. tristis, 502; multicaule, 477, 599; multijugum, 477, 599; multisectum, 480; musæfolium, 475; mysurense, (*Roth.*) 483, (*Wall.*) 484, 599; Nidus, 475, 599, var. Simonsiana, 475; nigripes, 490, 491, 494, 599, var. dissecta, 491, var. seleopteris, 491; niponicum, 493; nitens, 600; nitidum, 483, 599, 600; normale, 477, 599, var. minor, 477, var. opacum, 478; obliquum, 605; odontophyllum, 604; opacum, 477; ovatum, 599; oxyphyllum, 489, 491, 493, 600, 602, 603, var. kulhaitense, 494, var. oxyphyllum, 494; pachyphyllum, 476; parallelum, 500, 600; parvulum, 485, 604; pavoninum, 478; pectinatum, 491, 600; pekinense, 483; pellucidum, 600, 606; penangium, 599; phyllitidis, 475; pinnatifido-pinnatum, 497; planicaule, 479, 482, 599; polymorphum, 500, 501, 502, 503, 600; polyodon, 479; polypodioides, 501, 503, var. effusior, 501, var. sublatifolia, 501, var. typica, 501, var. vestita, 501; polyrrhizon, 499; porphyrocaulon, 481; porrectum, 498, 599, 600, 604; Prescottianum, 600; procerum, 427, 495, 501, 604, var. mishmica, 496; proliferum, 599, 600; prolongatum, 486; puberulum, 604; pubescens, 503; pulchellum, 483, 599; pulchrum, 604; pumilum, 482, var. hymenophylloides, 482; pusillum, 477; radiatum, 505, 597; resectum, 481;

- reticulatum, 505, 599; *Ruta*, 485; *rutæfolium*, 486; *Ruta-muraria*, 482; *Schkuhrii*, (*Hk.*) 498, 503, (*Mett.*) 499, (*Thw.*) 503; *septentrionale*, 478; *sepulchrale*, 483; *ser-ræforme*, 481; *sikkimense*, 427, 500, 501; *simplex*, 475; *soboliferum*, 498, 599; *sorzogonense*, (*Hk.*) 500, (*Presl*) 500, 600; *spec-tabile*, 490, 502, 600; *spinulosum*, 486, var. *subtriangularis*, 487; *splendens*, 483; *squarrosum*, 489; *stans*, 486; *stenophyllum*, 476; *Stoliczkæ*, 500, 604; *stramoneum*, 494; *strigillosum*, 490, 491; *sub-sinuatum*, 496, 599; *succulentum*, 427, 502; *sylvaticum*, (*Presl*) 497, 599, (*Bl.*) 500, 501; *Tavoyanum*, 480, 603; *tenerrimum*, 482; *tenui-folium*, 485, 599; *tenuifrons*, 490, 491, 599; *thelypteroides*, 488; *Thwaitesii*, 499; *tingens*, 476; *to-mentosum*, (*Hk.*) 498, 599, (*Mett.*) 499; *torrentium*, 427, 500; *tra-peziforme*, 481, 604; *Trichomanes*, 477, 599, var. *anceps*, 477; *um-brosum*, 496, 501, var. *procerum*, 495; *unilaterale*, 481, 599, var. *udum*, 481; *urophyllum*, 480, 599; *varians*, 483, 485, 486, 600; *viride*, 477; *Wichuræ*, 479; *Wightianum*, 604.
- Asteranthos*, 1, 17; analysis of, 21; *brasiliensis*, 18.
- Astrothelium ochrocleistum*, 278; *prostratum*, 278; *pyrenastroides*, 278.
- Athyrium allantodioides*, 488; *aspi-dioides*, 495; *Atkinsoni*, 487; *aus-trale*, 496, 502; *brevisorum*, 494; *Clarkei*, 490; *falcatum*, 487; *fimbriatum*, 494; *Griffithii*, 502; *gymnogrammoides*, 490, var. *ery-throrachis*, 491; *Hookerianum*, 486; *macrocarpum*, (*Bedd.*) 489, (*Fée*) 485; *multicaudatum*, 502; *nigripes*, 490; *oxyphyllum*, 493, 531; *rubricaulæ*, 492; *stramineum*, 493; *thelypteroides*, 488.
- Auricularini*, 126; of Brisbane, 403.
- Australian fungi, 399.
- Auxemma*, gen. char. of, 23; *Gard-neriana*, 24.
- Auxemneæ*, description of, 23.
- Avena elatior*, fruit of, 163; *sterilis*, torsion of awn, 157, 162.
- Awn, on the bending of, at the knees, 159; hygroscopic action of, in *Stipa*, 150, in *Avena*, 162.
- Azadirachta*, 234.
- Bactridium flavum*, 406.
- Baggiettoa ocellata*, 279.
- Baker, J. G., Report on the Liliaceæ, Iridaceæ, Hypoxidaceæ, and Hæ-modoraceæ of Welwitsch's Angolan Herbarium, 245.
- Ballia Brunonia*, struct. of, 216; *calli-tricha*, struct. and mode of growth, 211, cells of stems of, 215, 222, joints with stoppers, 212; *Hombro-niana*, 226; *Mariana*, 227; *Robertiana*, 227; *scoparia*, struct. of, 228.
- Barringtonia*, 55; *acuminata*, 102; *acutangula*, 79, (*Benth.*) 86, (*Bl.*) 75, (*Gaert.*) 80, (*Korth.*) 87, (*Roeb.*) 82, (*Span.*) 93, (*Thw.*) 81, (*W. & A.*) 83; *alata*, 70, 72, 93; *alba*, 65; *caffra*, 78; *ceylanica*, 77; *conoi-dea*, 65, 71, 72; *costata*, 88; *cylindrostachys*, 72; *cylindrostachya* (*Griffith*), 100; *edulis*, 76; *elon-gata*, 69; *excelsa*, (*Benth.*) 74, (*Bl.*) 74, 112, (*Gray*) 75; *Hors-fieldia*, 85; *insignis*, 75; *inter-media*, 73; *macrocarpa*, 109; *ma-crophylla*, 61; *Montrouzieri*, 89; *neocaledonica*, 103; *nitida*, 88; *obtusangula*, 81; *racemosa*, (*Bl.*) 67, 69, (*Gaud.*) 75, (*Griffith*) 71, (*Oliv.*) 78, (*Seem.*) 73, (*Thw.*) 77, var. *elongata* (*Bl.*) 109; *Reinward-tii*, 88; *reticulata*, 87; *rosca*, 102; *rubra*, 69; *samoënsis*, 74, 75; *sarcostachys*, 102; *serrata*, 87; *speciosa*, (*Forst.*) 56, (*Bl.*) 55, 58, (*Gaert.*) 68, (*Lin. fil.*) 61, (*W. & A.*) 63; *spicata*, 85; *sumatrana*, 103; *timoriensis*, (*Bl.*) 109, var. *alata* (*Miq.*) 93; *Vriesii*, 106.
- Barringtoniaceæ*, calyx of, 47; fruit of, 48; embryo of, 48; geograph. distrib. of, 54; inflorescence of, 47; ovary of, 48; petals of, 47; posi-tion of, 53; stamens of, 48; style of, 48; synop. char. genera of, 54.
- Belvisia* (and position of), 1, 17.
- Bennett, A. W., Prelim. Note no
- Growth of Female Flower-stalk: of *Vallisneria spiralis*, 133.
- Berkeley, Rev. M. J., and C. E. Broome, on Fungi of Brisbane, Queensland, 399.
- Blechnidium melanopus*, 475.
- Blechnopsis Cumingiana*, 474; *elon-gata*, 474; *latifolia*, 474; *nitida*, 474; *pyrophila*, 474; *salicifolia*, 474; *stenophylla*, 474.
- Blechnum brasiliense*, 474; *cartila-gineum*, 473; *elongatum*, 474; *Finlaysonianum*, 474, 604; *fla-bellatum*, 505; *glandulosum*, 596; *imbricatum*, 474; *melanopus*, 474; *moluccanum*, 474; *nitidum* (*Presl*), 474, var. *contractum* (*Hook.*) 474; *orientale*, 474, 596; *pteridioides*, 474; *pyrophyllum*, 474; *radiatum*, 505; *serrulatum*, 474; *striatum*, 474.
- Boragineæ, peculiarities infloresc. of, 615; self-fertilization sp. of, 375.
- Botrychium daucifolium*, (*Wall.*) 587, (*Hook.*) 596; *lanuginosum*, 588, 596; *Lunaria*, 587; *subbifolia-tum*, 587; *subcarnosum*, 587, 588, 596; *virginianum*, 588, 596; *zey-lanicum*, 587.
- Botryopteris mexicana*, 587.
- Botryopsis luzonensis*, 84.
- Bourreria clariuscula*, 26; *succulenta*, 26.
- Bovisca lilacina*, 406.
- Bowringia insignis*, 571.
- Bracts as adapted to self-fertilization, 380; uplifting of, 616.
- Brainea insignis*, 571.
- Brisbane, Queensland, fungi of, 399.
- British Lichens, new, 237.
- Broome, C. E., and Rev. M. J. Berke-ley, on Fungi of Brisbane, Queens-land, 399.
- Bulbine asphodeloides*, 261.
- Burdock, bur of, 165.
- Butonica alata*, 70; *alba*, 65; *apicu-lata*, 78; *caffra*, 78; *calyptrata*, 76; *ceylanica*, 77; *edulis*, 76; *incolta*, 71; *intermedia*, 73; *procera*, 74; *racemosa*, (*Juss.*) 66, (*Bl.*) 67; *rosata*, 72; *rubra*, 70; *Rumphiana*, 67, 68; *samoënsis*, 75; *splendida*, 60; *terrestris*, 69; *terrestris rubra*, 69; *Tsjeria Saamstravadi*, 70.

- Butonjca*, embryo of, 50, 52, 53; hist. and charac. of, 65.  
*Butonicoides crenata*, 90, 91.  
*Byblis gigantea*, striated cells of, 166.  
*Cabralea*, 236.  
*Cæomacei* (of Brisbane), 407.  
*Calicium kylemoriense*, 242.  
*Callipteris esculenta*, 503.  
*Calocera viscosa*, 127.  
 Calycanthemy of Primulaceæ, 286.  
 Calyx as adapted to self-fertilization, 380.  
*Camellia japonica*, blight on leaves of, 301; minute structure of, 303.  
 Campanulaceæ, self-fertilization sp. of, 368.  
*Campteria anamallayensis*, 469; *bi-aurita*, 469; *Rottleriana*, 469; *Wallichiana*, 470.  
*Campylandra dachystachys* (*Welw.*), 256.  
*Carapa procera*, 235.  
*Cardiochlena alata*, 539.  
*Careya arborea*, 97, var. *australis*, 91; *crenata*, 90, 91; *herbacea*, 96, 97; *macrostachya*, 85, 104; *orbiculata*, 98; *pendula*, 96, 99; *spherica*, 66, 97.  
*Careya*, charac. of, 96; history of, 95; seed of, 48.  
*Caryocar*, embryo of, 51.  
 Caryophyllaceæ, self-fertilization sp. of, 353, 392.  
*Cedrela Glaziovii*, 236; *sinensis*, distribution of, 233.  
*Cenangium lichenoideum*, 404.  
*Cephalomanes auriculatum*, 441; *javanicum*, 442; *rhomboideum*, 442; *Zollingeri*, 442.  
 Ceratophylleæ, self-fertilization sp. of, 393.  
*Ceratopteris thalictroides*, 471, 597.  
*Ceterach aureum*, 504; *officinatum*, 504; *pedunculata*, 570.  
*Chaetocladium*, fructification of, 420.  
*Chasalia coffeoides*, 176; *divaricata*, 176; *Fontanesii*, 175.  
*Cheilanthes albomarginata*, 427, 456, 457; *argentea*, 458, var. *sulphurea*, 458; *bullosa*, 457; *chrysophylla*, 458; *contigua*, 459, 597; *Dalhousiæ*, 457; *dealbata*, 457, 596; *farinosa*, 456, 457, 604, var. *chrysophylla*, 458, var. *concolor*, 457, var. *Dalhousiæ*, 456, 457, var. *vestita*, 457; *fimbriata*, 455; *fragrans*, 454; *lucida*, 459, 596; *micrantha*, 455, 596; *mysurensis*, 596; *nitidula*, 460; *odora*, 454; *pallens*, 596; *pallida*, 546; *rigidula*, 457, 604; *rufa*, 456, 457; *rupestris*, 455, 596; *setigera*, 528; *Sieberi*, 455; *suaevolens* (*Swz.*), 454; *subvillosa*, 456; *Szovitzii*, 454, 455; *tenuifolia*, 455, 596; *tomentosa*, 457, 597; *varians*, 455, 597.  
*Cheiranthus Cheiri*, æstivation of, 190.  
 Chenopodiaceæ, self-fertilization sp. of, 377, 393.  
*Chetobromus Dregeanus*, awn of, 164.  
*Chiodecton inconspicuum*, 280; *conchyliatum*, 280; *moniliatum*, 280; *sinuosum*, 280.  
*Chisocheton paucijugum*, 236.  
*Chlorophytum andongense*, 260; *bracteosum*, 260; *debile*, 260; *filipendulum*, 260; *lanceifolium*, 260; *leucolepis* (*Welw.*), 259; *macrophyllum*, 259; *stenopetalum*, 260.  
*Chnophora glauca*, 432.  
 Choanephora, as illustrating conidial fructification in the Mucorini, by Dr. Cunningham, 409; Brefeld's researches, 419; chlamydo-spores of, 428; conclusions from phenomena of develop., 419; conidia, devel. of, 414; exper. artificial cultivation of, 413; formation of zygospore of, 412; germinal tubes of, 414; history of, 409; minute structure of, 410; reproductive bodies differ, 418; Van Tieghem and Le Monnier's researches, 420; variations from nourishment of, 416.  
*Chrysodium aureum*, 582; *cayennense*, 582; *D'Urvillei*, 582; *fraxinifolium*, 582; *sculpturatum*, 582.  
*Chydenanthus*, charac. of, 111; *excelsus*, 112, hist. of, 112.  
*Cibotium assamicum*, 435; *Barometz*, 435; *glaucoscens*, 435.  
*Circaea*, 178.  
*Circinella umbellata*, 406.  
*Cladosporium herbarum*, 407.  
 Clarke, C. B., Review of the Ferns of Northern India, 425.  
*Clavaria argillacea*, 403; *rugosa*, 403; *stricta*, 403.  
*Clavariæ*, 127; of Brisbane, 403.  
 Clusiaceæ, embryo in, 50.  
*Cœnopteris appendiculata*, 485; *furcata*, 600; *vivipara*, 600.  
*Coffea*, African species of, by W. P. Hiern, 169.  
*Coffea Afzeliï*, 169, 174; *arabica*, (*L.*) 169, 170, (*Benth.*) 171, 172, (*β, Willd.*) 173, var. *leucocarpa*, 171; *bengalensis*, 169; *brevipipes*, 169; *capitata*, 176; *cymosa*, 175; *divaricata*, 176; *hirsutus*, 176; *hypoglauca*, 169, 173; *jasmoides*, 169, 175; key to classification, 170; *Kraussiana*, 176; *laurifolia*, 170; *laurina*, 175; *liberica*, 169, 171; *macrocarpa*, 169, 173; *mauritanica*, 169, 173; *melanocarpa*, 169, 173; *microcarpa*, 164, 174; *mozambicana*, 175; *Perrotettii?* (*Steud.*), 174; *raccmosa*, 169, (*Lour.*) 175; *ramosa* (*Roem. et Schult.*), 175; *rupestris*, 169, 174; *stenophylla*, 169, 172; *subcordata*, 169, 174; *sylvestris*, 173; *tenuiflora*, 169; *tetrandra*, 169; *zangoobarica*, 169, 172.  
*Coleochæte scutata*, 311.  
 Commersonia, 68.  
 Compositæ, plumes on seed of, 165; self-fertilization sp. of, 367, 392.  
 Conidial fructification in the Mucorini, by Dr. Cunningham, 409.  
 Contabescence, 381.  
 Contribution to the Lichenographia of New Zealand, by Charles Knight, 275.  
 Convolvulaceæ, self-fertilization sp. of, 368, 392.  
*Corallina monile*, 204; *officinalis*, 203, histology, frond of, 204; *opuntia*, 202, Decaisne's obs. on, 198; *tridens*, 200, 202; *tuna*, 200, 202; *vermicula*, 200.  
 Corallinaceæ, brilliant colours of, 200; colouring-matter of, 200; Duncan and Nelson on, 197, 200; of Bermuda, struct. of, 200.  
 Corolla as adapted to self-fertiliza-

- tion, 380; irregular plants showing, 184; regular plants showing, 184.
- Corolla-lobes, develop. after stamens in *Arenaria trinervis*, 196; in *Cerastium glomeratum*, 196; in *Lychnis dioica*, 196; in *Ranunculus acris*, 196; in *Stellaria Holostea*, 196; in *Veronica Chamædrys*, 196.
- Corticium læve, 403; lævigatum, 127; Mougeotii, 127; violaceo-lividum, 127.
- Coussarea (*Aubl.*), 169.
- Crassulaceæ, modification cymes of, 617.
- Craterispermum laurinum, 175.
- CreMASpora africana, 176.
- Cruciferae, binary symmetry of, 193; marginal placentas of, 192; self-fertilization sp. of, 350, 391.
- Cruciferous flower, structure of, Rev. G. Henslow on, 177.
- Cryptogramme acrostichoides, 459; Brunoniana, 459, 603; crispa, 459, 460, 603, var. acrostichoides, 460.
- Cunningham, Dr. D. Douglas, on the occurrence of Conidial fructification in the Mucorini, illus. by Choanephora, 409; on Mycoidea parasitica, and the part it plays in formation of certain Lichens, 301.
- Curculigo gallabatensis, 266.
- Currey, Fred., on a collection of Fungi made by Mr. Sulpiz Kurz, 119.
- Cyathea barbata, 435; bipinnata, 605; Brunonis, 599; excelsa, 599; robusta, 599; spinulosa, (*Wall.*) 429, 430, 431, 599, (*Bedd.*) 429, (*J. Scott*) 430; venulosa, 433, 599.
- Cyathus campanulatus, 405; fime-tarius, 405; intermedius, 128.
- Cyclamen, double petals, 287.
- Cyme, dichotomous, 618; scorpioid, origin of, by Rev. G. Henslow, 613, 618; spicate, 618.
- Cympolia barbata, 198.
- Cyrtomium (subgen.), 512; caducum, 512; caryotideum, 512; falcatum, 512.
- Cystopteris dimidiata, 444; fragilis (*Bernh.*), 450, f. Huteri (*Milde*), 450; nodosa, 444; retusa, 450, 492; setosa, 450; spinulosa, 486; squamata, 444.
- Dædalea discolor, 125; Sprucei, 402; tenuis, 125; unicolor, 403; zonata, 125.
- Danthonia, awn of, 164.
- Darwin, Francis, on the Hygroscopic Mechanism by which certain Seeds are enabled to bury themselves in the ground, 149.
- Dasystachys, 255; campanulata, 256; colubrina, 256; falcata, 256; pleio-stachya, 255.
- Davallia achilleifolia, 600; affinis, 444; alpina, 442; angustata, 600; assamica, 443; Boryana, 451; bulbata, 445; caudata, 605; chærophylla, 444; chinensis, 449, 600; Clarkei (*Hk. & Bk.*), 443; conii-folia, 600; dareæformis, 427, 443; dimidiata, 445; divaricata, 445; elegans, 600; flaccida, 447, 448, 605, var. pilosula, 449, var. pubera, 449, var. pyramidata, 449; flagellifera, 600; Griffithiana, 445; hemiptera, 451; hirta, 447; Hookeriana, 446, 447, 605; immersa, 443; jamaicensis, 448; khasiyana, 447; Kurzii, 446; ligulata, 444; lobulosa, 600; lonchitidea, 446, 600; marginalis, 446, 600, 605, var. calvescens, 447; membranulosa, 442, 443; micans, 443; moluccana, 605; multidentata, 442, 443, 602; nodosa, 434, 443, 444, 602; ornata, 600; parallela, 600; parvula, 600; pedata, 442, 600; pilosa, 448; pilosula, 447; platyphylla, 446, 600; polyantha, 445; polypodioides, 447, (*Don*) var., 605, var. hirta, 448, var. pilosula, 448, var. rhomboidea, 448, var. strigosa, 448; proxima, 447; puberula, 448; pulchra, 444, 445, var. pseudo-cystopteris, 444, 450; pyramidata, 448, 449; repens, 427, 451; rhomboidea, 447; Roxburghii, 447, 605; serriformis, 600; setosa, 450; scabra, 446, 447; schizophylla, 451; Speluncae, 448; sphæropteroides, 445; stipellata, 444; strigosa, (*Swartz*) 436, 447, (*Hk. & Bk.*) 447; subimbricata, 442; tenni-folia, 449, 600; trichomanoides, 451; trichosticha, 448; urophylla, 446, 447, 605; villosa, 446, 600.
- DeCandolle, Casimir, on the geographical distribution of the Meliaceæ, 233.
- Dematiei (of Brisbane), 407.
- Dennstaedtia deltoidea, 436.
- Diacalpa aspidioides, 434, 445, 603; fœniculacea, 427, 434; Hookeriana, 434; sphæropteroides, 455.
- Dialysis, rare cases of, 193.
- Diatrype rugosa, 130.
- Dichasium of Schimper, 618, 619.
- Dicksonia appendiculata, 436, 596; arborescens, 596; assamica, 435; Barometz, 435; deltoidea, 436; Elwesii, 436; Griffithiana, 435; moluccana, 604; polypodioides, 447; scabra, 436, 604.
- Dictyocline Griffithii, 576.
- Dictyopteris (subgen.), 548.
- Didymium farinaceum, 406.
- Didymoglossum euphlebium, 440; Filicula, 440; late-alatum, 440; plicatum, 440, racemosum, 440.
- Digrammaria ambigua, 503.
- Dipcadi comosum, 247; lateritium, 247; oxylobum, 246; Welwitschii, 246.
- Diplazium (subgen.), 496; alternifolium, 497; bantamense, 497; bulbiferum, 498, 499; dilatatum, 502; esculentum, 503; falcatum, 497, 569; fraxineum, 569; fraxinifolium, 497; Griffithii, 502; Jerdoni, 502; lanceum, 496; lasiopteris, 499; lobulosum, 478; longifolium, 478; malabaricum, 503; malaccense, 499; pinnatifido-pinnatum, 497; polypodioides, 501; Schkuhrii, 499, 503; sorzogonense, 500; speciosum, 499; Stoliczkae, 500; sylvaticum, 498; thelypteroides, 488; Thwaitesii, 499; tomentosum, 498.
- Dipteris (subgen.), 555; Wallichii, 555.
- Doryopteris sagittifolia, 471.
- Double primroses, 287, 293.

- Doxomma* (n. gen. *Miers*), 98, 99; *acuminatum*, 102; *angustatum*, 105; *cochinchinense*, 101; *cylindrostachya*, 100; *macrostachyum*, 104; *magnificum*, 105; *neocaledonica*, 103; *pendulum*, 99; *rigidum*, 104; *sarcostachys*, 102, 103; *sumatrana*, 103; *Vriesii*, 106.
- Dracæna* *acaulis*, 252; *fragrans*, 252; *interrupta*, 252; *monostachya*, 252; *nitens*, 252; *parviflora*, 252.
- Drymoglossum* *Beddomei*, 576, 598; *carnosum*, 575, 598; *piloselloides*, 575, 598; *subcordatum*, 575.
- Drynaria* (subgen.), 555; *coronans*, 557, 582; *mollis*, 556; *propinqua*, 556; *quercifolia*, 556, 582.
- Dysoxylum* *spectabile*, distrib. of, 233.
- Elaphoglossum* *conforme*, 576; *laurifolium*, 576; *viscosum*, 577.
- Ella-medella-gass* = *Stravadium* *acutangulum*, 80.
- Embryo in *Clusiaceæ*, 50.
- Equisetum* *arvense*, 594; *bullata*, 600; *chærophylla*, 600; *debile*, 594, 603; *diffusum*, 594, 600; *elongatum*, 595, 600; *flaccida*, 600, and var., 600; *immersa*, 600; *laxum*, 595; *membranulosa*, 600; *nodosa*, 600; *pallens*, 595, 603; *palustre*, 594; *pilosa*, 600; *polypodioides* (*Don*), var., 600; *puberula*, 600; *pulchra*, 600; *pyramidata*, 600; *ramosissimum*, 595, *ramosum*, 595; *rhomboides*, 600; *robustum*, 595; *scirpioides*, 600; *scoparium*, 594, 603; *stipellata*, 600; *virens*, 600; *virgatum* (*Bl.*), 595.
- Eremobryoid series of *Polypodium*, 549.
- Erineum, tubes of, 166.
- Eriospermum* *andongense*, 261; *flexuosum*, 261; *ophioglossoides*, 262; *paludosum*, 261; *stenophyllum*, 261.
- Erodium*, manner in which seed of, is buried, 155.
- Eugenia* *acutangula*, (*Linn.*) 70, 79, 80, (*Lour.*) 101; *crenata* (*Soland.*), 91; *racemosa*, (*Linn.*) 66, (*Seem.*) 73, (*Forst.*) 73, (*Roxb.*), 82.
- Eunephrodium* (subgen.), 529, 547.
- Eupapaveræ*, 180.
- Euphorbiaceæ*, self-fertilization sp. of, 378, 393.
- Eupolypodium* (subgen.), 549.
- Forctia* *apodanthera*, 169.
- Ferns, difficulties with regard to nomenclature, 426.
- Ferns of N. India, review of, 425.
- , change of names, 427; Col. *Beddome's* tab. of distrib. of species, 426; new species, 427; *Wallich's* collection, how to be quoted, 427, tabular reduction of, 595.
- Fissurina* *novæ-Zelandiæ*, 281.
- Flindersia*, 235; *Fournieri*, 236.
- Floral æstivations, diagrams of, 177; introductory remarks on, 177; origin of, *Rev. G. Henslow* on, 177.
- Forms of æstivation, 178; causes which determine, 187, in irregular flowers, 188, in regular flowers, 187; cochlear, cochleate (footnote), 182; convolute and contorted, 179, 185; degrees of frequency, 179; distichous equitant, 178, 180; equitant, 180; half-equant, 178, 180; half-imbricate, 178, 182; imbricate proper, 178, 184; imbricative, 179; induplicative, 179; numerical order of parts of whorls, 186; pentastichous, 178, 182; quincuncial, 178, 182; reduplicative, 179; reversal of the spiral, 186; straight or open, 179; tristichous, 178, 181; valvate, 179, 185; valvative, 180; varieties in same species of diff. genera, 189.
- Fuchsia* *coccinea*, æstivation of, 190.
- Fumago* *salicina*, 130.
- Fungi, on a collection of, made by *Mr. Sulpiz Kurz*, 119.
- Galanthus* *nivalis*, æstivation of, 190.
- Galium* *Aparine*, bur of, 165.
- Geaster* *saccatus*, 406; *striatus*, 406.
- Gentianaceæ*, self-fertilization sp. of, 368.
- Geographical distribution of the *Meliaceæ*, 233; map of (Pl. 31), 236; summary of, 236; tabular arrang. of (Pl. 30), 236.
- Geraniaceæ*, burying powers of, 165; self-fertilization sp. of, 357, 392.
- Gethyllis* *pilosa*, 267.
- Geum*, bur of, 165.
- Gladiolus* *andongensis*, 269; *angolensis*, 269; *bengnellensis*, 268; *brevicaulis*, 267; *cærulescens*, 267; *gregarius*, 268; *laxiflorus*, 268; *luridus*, 267; *multiflorus*, 269; *quartinianus*, 269; *splendens* (*Welw.*), 268; *Welwitschii*, 268.
- Glaux*, corolla wanting of, 287.
- Gleichenia* *arachnoides*, 428; *attenuata*, 598; *Bancroftii*, 428; *crassifolia*, 428; *dichotoma*, 428; *excelsa*, 428; *ferruginea*, 428; *gigantica*, 428, 598; *glauca*, 428, 598; *Hermanni*, 428, 598; *japonica*, 428; *Klotzschii*, 428; *lanigera*, 428; *linearis*, 428, 598; *longissima*, 428; *pteridifolia*, 428.
- Gloriosa* *superba*, 262; *virescens*, 262.
- Goniophlebium* (subgen.), 550; *amœnum*, 550; *argutum*, 551; *erythrocarpum*, 565; *Hendersoni*, 550; *lachnopus*, 551; *microrhizoma*, 551; *serratifolium*, 552; *subauriculatum*, 552.
- Goniopteris* (subgen.), 547; *lineata*, 547; *multilineata*, 548; *penangiana*, 548; *prolifera*, 548.
- Gramineæ*, self-fertilization sp. of, 379, 393.
- Grammitis* *acuminata*, 570, 595; *affinis*, 569, 595; *arborescens*, 568; *aurita*, 568; *caudata*, 569, 595; *Ceterach*, 504; *decurrens*, 571, 595; *diversifolia*, 595; *scopendrina*, 570; *cuspidata*, 570; *Finlaysonia*, 571, 603; *flavoscens*, 570, 595; *Hamiltoniana*, 570, 595; *involuta*, 570; *lanceolata*, 570; *macrophylla*, 570, 595; *microphylla*, 569; *procera*, 569, 595; *Totta*, 567, 568; *vestita*, 568, 595.
- Grandinia* *granulata*, 126.
- Graphiola* *phœnicis*, 130.
- Growth, curves of, *A. W. Bennett* on, 136; diagrams of, 137, 141; in *Ballia* *callitricha*, 211; in flower-stalk of *Hyacinth*, 139; in fl.-stalk of *Vallisneria*, 133.
- Guarea* *filiformis*, 236; *trichilioides*, 236.

- Guepinia ramosa*, 127; *spathularia*, 403.
- Gustavia alata*, 93; *globosa*, 86; *valida*, 93, 94.
- Gymnogramme Andersoni*, 568; *aspidioides*, 567, 569; *aurita*, 544, 568, var. *Levingii*, 568; *decurrens*, 570; *elliptica*, 570, 595, 603; *fraxinea*, 569, 595, var. *pilosa*, 569; *Hamiltoniana*, 570, 595; *involuta*, 570, 595; *javanica*, 569; *lanccolata*, 570; *Lovei*, 567; *Marantæ*, 567; *microphylla*, 569; *obtusata*, 568; *opaca*, 568, 600; *serrulata*, 569; *stegnogramme*, 569; *Totta*, 567; *vestita*, 568, 595.
- Gymnopteris axillaris*, 579; *decurrens*, 579; *Féei*, 579; *lanccolata*, 579; *minor*, 579; *normalis*, 579; *spicata*, 582; *tricuspis*, 581; *variabilis*, 579.
- Gymnosphaera gigantea*, 496; *glabra*, 433.
- Hamodoraceæ*, Angolan, 245.
- Haloragiaceæ*, self-fertilization sp. of, 392.
- Harmena*, 53.
- Haworthia angolensis*, 263.
- Heliotrope*, scorpioid racemes of, 616.
- Helminthosporium Ravellinii*, 407.
- Helminthostachys dulcis*, 595; *zeylanica*, 587, 595.
- Holotium citrinum*, 128.
- Helvella californica*, 423.
- Helvellacei*, 128; of Brisbane, 404.
- Hemidictyon Finlaysonianum*, 504.
- Hemidictyum* (subgen.), 504; *Brunonis*, 505; *Ceterach*, 504.
- Hemionitis arifolia*, 576, 595, 604; *Boryana*, 573; *cordata*, 576; *cordifolia*, 576, 595; *coriacea*, 572; *esculenta*, 503; *Griffithii*, 576; *hastata*, 575, 604; *opaca*, 568; *pothifolia*, 571; *reticulata*, 573; *sagittata*, 575; *Wilfordii*, 575.
- Hemitelia Beddomi*, 429; *Brunoniana*, 427, 430, 431, var. ? *Scottii*, 431; *decipiens*, 430; *latebrosa*, 432.
- Henslow, Rev. George, on Origin of prevailing systems of Phyllotaxis, 37; on Origin of so-called Scorpioid Cyme, 613; on the Origin of Floral Æstivations, 177; on the Self-fertilization of Plants, 317.
- Hermannia*, whorls of, 177.
- Heteroneuron diversifolium*, 580; *heteroclitum*, 580; *proliferum*, 581.
- Heteropogon contortus*, awn of, 164; *melanocarpus*, awn of, 164.
- Hexagonia Kurzi*, 126; *polygramma*, 125, 403; *tenuis*, 125.
- Heynea*, 264.
- Hiern, W. P., on the African species of the genus *Coffea*, Linn., 169.
- Hinneola auricula Canis*, 128; *auricula Judæ*, 128, 404; *polytricha*, 404.
- Histology of Corallineæ, 197; DeCaisne on, 197; Quekett on, 197; Rosanoff on, 199.
- Holcus*, awn of, 164.
- Humata pedata*, 442.
- Hyacinth*, on growth of flower-stalk of, 139, 143.
- Hydnei*, 126; of Brisbane, 403.
- Hydnum auriculatum*, 126; *gelatinosum*, 126; *udum*, 126.
- Hydrocharis*, whorl flower of, 177.
- Hydroglossum auriculatum*, 584; *circinatum*, 583; *dichotomum*, 583; *flexuosum*, 584; *japonicum*, 585; *longifolium*, 583; *pedatum*, 583; *pinnatifidum*, 584.
- Hygrophorus miniatus*, 399.
- Hygroscope*, how made, 155.
- Hygroscopic mechanism* by which certain seeds are enabled to bury themselves in the ground, 149; *hygroscopic torsion*, 166.
- Hymenesthes*, hist. and charac. of, 26; *nitida*, 26.
- Hymenochaete Cacao*, 403; *tenuissima*, 403.
- Hymenolepis ophioglossoides*, 582; *revoluta*, 582; *spicata*, 582.
- Hymenophyllum abietinum*, 437; *alatum*, 440; *badium*, 437, 439, 599; *Blumeianum*, 437; *Boryanum*, 598; *ciliatum*, 437, 439; *crispatum*, 437, 598; *dædaloum*, 437; *densum*, 436, 440, 598; *denticulatum*, 438, 439, var. *flaccidum*, 438, 439; *erosum*, 437; *exsertum*, 436, 437, 598; *Filicula*, 440; *fimbriatum*, 437; *flaccidum*, 439; *flexuosum*, 437; *himalaianum*, 437; *integrum*, 437; *javanicum*, 437, 439, 598, 599, var. *badium*, 438; *khasianum*, 438; *lave*, 604; *Levingii*, 427, 439; *macroglossum*, 436; *micranthum*, 437; *microsorium*, 437; *osmundioides*, 437; *polyanthos*, 437, 598, 599, var.  $\beta$ . *minor*, 437; *protrusum*, 437; *pycnocarpum*, 437; *Reinwardtii*, 437; *serpens*, 437, 599; *Simonsianum*, 438; *Smithii*, 439; *sphaerocarpum*, 437; *Telfairianum*, 598; *tenuis*, 605.
- Hyoscyamus*, floral axis of, 615.
- Hypematium onustum*, 525; *Ruppelianum*, 525.
- Hypochaete debilis*, 595.
- Hypocrea variabilis*, 130.
- Hypolepis hostilis*, 546; *pteridioides*, 468; *tenuifolia*, 546.
- Hypoxidaceæ*, Report on, of Welwitsch's Angolan herbarium, 245, 265.
- Hypoxis angolensis*, 266; *angustifolia*, 265; *canaliculata*, 265; *cuanzensis*, 265; *monanthos*, 266; *polystachya*, 266, var. *andongensis*, 266.
- Hypoxylon angolense*, 405; *octarioides*, 405; *concentricum*, 130, 405; *cretaceum*, 405; *marginalatum*, 130; *suborbiculare*, 130.
- Ileodictyon gracile*, 405.
- Incompletæ*, self-fertilization sp. of, 377.
- India proper, Kurz's divisions of, 425.
- India, Review of the Ferns of Northern, 425.
- Inflorescence*, steps towards scorpioid cyme, 617.
- Iridaceæ* (Angolan), 267; Report on, of Welwitsch's herbarium, 245.
- Iridopsis textilis*, 270.
- Irish lichens, new, 241.
- Irpex flavus*, 126, 403; *pallescens*, 126; *sinuosus*, 126; *zonatus*, 403.
- Jambos sylvestris Saamstravadi*, 66.



- Juncaceæ, self-fertilization sp. of, 379, 393.
- Kaulfussia æsculifolia, 585; assamica, 585; Korthalsii, 585; Lobbiana, 585.
- Knight, Chas., Contribution to the Lichenographia of New Zealand, 275.
- Kniphofia andogensis, 246; benguelensis, 246.
- Kraussia floribunda, 176.
- Kurz, Mr. Sulpiz, on a collection of Fungi made by, 119.
- Labiatae, self-fertilization sp. of, 374, 392.
- Lacaussadea appendiculata, 578; montana, 578; rhizophylla, 578.
- Lachnocladium furcellatum, 403.
- Lagerströmia, oscillating arrang. leaves of, (ftnote) 45, 613.
- Lagurus ovatus, awn of, 164.
- Lansium, 235.
- Lapeyrousia abyssinica, 272; cyaneascens, 272; fragrans, 272; littoralis, 272; odoratissima, 273.
- Laschia Thwaitesii, 403.
- Lastrea (subgen.), 512; angustifrons, 527; apiciflora, 518; aristata, 511; Atkinsoni, 526; barbigeræ, 522; bella, 525; Bergiana, 515; Blumei, 526; Boryana, 527; Brunoniana, 522; cana, 516; cochleata, 522, 582; conifolia, 511; erenata, 525; eburnea, 493; elongata, 513, 521, 523; Elwesii, 516; eriocarpa, 525; Fairbanki, 517; falciloba, 515; Falconeri, 522; Filix-Mas, 582, var. Clarkei, 518, var. odontoloma, 521; flaccida, 514, 528; fœniculacea, 434; gracilescens, 514; gracilis, 508; Hendersoni, 526; hirtipes, 513; immersa, 514; intermedia, 520; macrocarpa, 489; nigrescens, 515; oethodes, 516; patentissima, 520; platypus, 528; pulvinulifera, 435, 525; recedens, 526; sericea, 515; setigera, 528; setosa, 450; sparsa, 494, 524, var. obtusissima, 524; spectabilis, 519; spinulosa, 523; splendens, 527; Thelypteris, 517; tyloides, 516.
- Lathræa Squamaria, arrang. inflores. of, 613.
- Leaf, peculiar change of position of, 616.
- Lecanora jejuna, *Nyl.*, = *L. subdiluta*, *Leight.*, 146.
- Lecidca abietina, 278; advenula, 146; antrophila, 242; atro-morio, 276; Dilleniana, 278; endomelana, 239; formosa, 278; glaucomaria, 238; grumosa, 242; imponens, 238; impressula, 237; littoralis, 275; mciospora, 275; nigrescens, 276; parellaria, 238; petræa (*Flot.*), var. neo-zelandica, 277, var. violacea, 277; polospora, 241; premnea, 278; prosochoides, 146; ryssolea, 237; schistacea, 276; stellulata, 277; subargillacea, 275; subbadiotatra, 276; subcoarctata, 276; subdiluta, 145; subfarinosa, 277; subglobulata, 275; sublapicida, 276; subtubulata, 277; tephrizans, 237; tubulata, 277; Whakatipæ, 277.
- Leguminosæ, self-fertilization sp. of, 360, 392.
- Leighton, Rev. W. A., new British Lichens, 145, 237; new Irish Lichens, 241.
- Lentinus cæspitosus, 120; capronatus, 119; cyathus, 399; descendens, 119, 400; Dunalii, 400; exilis, 120, 400; fasciatus, 399; furfurosus, 120; glabratus, 120; inquinans, 120; irregularis, 121; Kurzianus, 120; Lecomtei, 400; Sajor Caju, 120; subdulcis, 400; velutinus, 120.
- Lonchitis albida, 121; betulina, 400; deplanata, 400; Palisoti, 121.
- Lepiota, 119.
- Leptochilus axillaris, 579; decurrens, 579; lanceolatus, 579; minor, 579.
- Leptostegia lucida, 459.
- Leucostegia immersa, 443.
- Lichenographia of New Zealand, contributions to the, 275.
- Lichens, new British, 145, 237; new Irish, 241.
- Liliaceæ, Report on, of Welwitsch's Angolan herbarium, 245.
- , self-fertilization sp. of, 357.
- Lindsaya attenuata, 450, 451, 598, 604; chinensis, 449; cultrata, 450, 451, 452, 598; decomposita, 593; dentata, 598; ensifolia, 452, 604; Finlaysonia, 452, 604; fabelulata, 451, 598; gracilis, 450; Griffithiana, 452; heterophylla, 452; interrupta, 451, 604; lanceolata, 452; lanuginosa, 541, 598; lobata, 598; Lobbiana, 450; longipinna, 452, 634; lucida, 450, 598; neriiforme, 542; oblongifolia, 451; odorata, 450; pallens, 450, 598; parasitica, 604; pectinata, 451; pentaphylla, 452; polymorpha, 451, 598; pteroides, 452, 604; recurvata, 598; repens, 427, 451; scandens, 451; serpens, 598; striata, 451; tenera, 451; tenuifolia, 449.
- List of Fungi from Brisbane, Queensland, with descrip. of new Species, by the Rev. M. J. Berkeley and C. E. Broome, 399.
- Litobrochia aurita, 471; ludens, 470; pedata, 470.
- Lomaria adnata, 472; articulata, 473; attenuata, 596; aurea, 458; 596; Boryana, 599; callosa, 472; carnifolia, 458, 596; euphlebia, 472, 473, 596; glauca, 472; limonifolia, 577, 596; Patersoni, 472; pyenophylla, 472, var. 472, 474; scandens, 577, 596; secunda, 596; semicordata, 472, 473; serpens, 579, 596; spondiæfolia, 596; Wightiana, 604.
- Lonchitis hirsuta, 604.
- Loxogramme involuta, 570; lanceolata, 570.
- Lycoperdacci, 128.
- Lycoperdon brasiliense, 128; gemmatum, 128; pusillum, 128, 406.
- Lycopodium aloifolium, 590, 598; annotinum, 592, 593; argenteum, 598; atroviride, 597; bryopteris, 604; cæspitosum, 594; casuarinoides, 593; caulescens, 598; cernuum, 591, 598, var. sikkimensis, 592; ceylanicum, 590; circinale, 604; clavatum, 592, 598; comans, 593; complanatum, 593, 604; concinnum, 597; curvatum, (*Dalz.*)

- 594, (*Switz.*) 591; *divaricatum*, (*Hk. & Grev.*) 594, (*Wall.*) 592, 598; *elegans*, 598; *empetrifolium*, 590; *epicæfolium*, 591; *flicale*, 593; *flexuosum*, 604; *fulcratum*, 598; *gnidioides*, 598; *gramineum*, 590; *Hamiltonii*, 590, 598, var. *petiolata*, 590; *Heyneanum*, 592, 598; *Hookeri*, 591, 597; *javanicum*, 591; *lævigatum*, 604; *lucidulum*, 590; *miniatosporum*, 594; *mirabile*, 593; *myrtifolia*, 593; *obtusifolium*, 590, 598; *ornithopodioides*, 604; *pallidum*, 606; *Phlegmaria*, 592, 598, var. *nummularifolia*, 593; *proliferum*, 591; *protensum*, 591; *pubescens*, 598; *pulcherrimum*, 590, 597; *pulvinatum*, 606; *rotundifolium*, 593, 604; *rubellum*, 593; *sabinæfolium*,\* 593; *semicordatum*, 598; *serratum*, 591, 597; *setaccum*, 590, 591, 597, var. *pulcherrima*, 590, var. *subulifolia*, 590; *squarrosum*, 591, 597, 606; *subdiaphanum*, 598; *subulifolium*, 590, 597; *sulcinervium*, 591; *tetragonostachyum*, 598; *thyoides*, 593; *trichiatum*, 592; *ulcifolium*, 591; *urostachyum*, 597; *vernicosum*, 589, 590; *verticillatum*, 591, 597, 606; *vulcanicum*, 592; *Wightianum*, 593, 604; *Willdenovii*, 598.
- Lygodium circinatum*, 583, 599; *dichotomum*, 583, 599; *Finlaysonianum*, 585, 604; *flexuosum*, 584, 585, 599; *japonicum*, 584, 604; *longifolium*, 583, 584, 599; *microphyllum*, 583, 584, 599; *pedatum*, 583; *pinnatifidum*, 584; *polystachyum*, 584, 599; *pubescens*, 584, 604; *scandens*, 584; *semibipinnatum*, 585; *semihastatum*, 584; *serulatum*, 584; *tenuis*, 585.
- Lysimachia nummularia*, 287.
- Lythraceæ*, self-fertilization sp. of, 363, 392.
- Macrochloa arenaria*, awn of, 164.
- Malvaceæ*, self-fertilization sp. of, 356, 392.
- Mammea asiatica*, 58, 61.
- Marasmins*, 119; *ramealis*, 119.
- Marattia fraxinea*, 599.
- Masters, M. T., on some points in the Morphology of the Primulaceæ, 285.
- Matonia pectinata*, 599.
- Mechanism of torsion, 157.
- Medella* = *Stravadium acutangulum*, 80.
- Megadendron*, 109; *ambiguum*, 110; *macrocarpum*, 109; *pallidum*, 110.
- Melaspilea amphorodes*, 281; *metabola*, 281; *vermifera*, 146.
- Melia Azedarach*, distrib. of, 233, 235.
- Meliaceæ, on the geographical distribution of, 233.
- Melobesiae, filamentous processes of, 207; Rosanoff on histology of, 199, 207.
- Menichea rosata*, 72.
- Meniscium Cumingii*, 571; *cuspidatum*, 547, 571, 572, 596, var. *longifrons*, 572, 596; *deltigerum*, 572, 580, 596; *erosum*, 571, 596; *longifrons*, 596; *Parishii*, 571; *proliferum*, 548; *reticulatum*, 572; *salicifolium*, 596; *simplex*, 571; *triphylum*, 571, 596; *villosum*, 571.
- Mertensia dichotoma*, 428; *emarginata*, 428; *glabra*, 428; *glauca*, 428.
- Mesocarpus nummuloides*, (fnote) structure of, 212.
- Meteorus coccineus*, 83.
- Microgonium bimarginatum*, 440; *sublimbatum*, 439.
- Microlepidia caudigera*, 446; *hirta*, 447; *platyphylla*, 446; *pteropus*, 443; *scabra*, 447; *spelunca*, 448; *strigosa*, 436, 447; *urophylla*, 446.
- Micropeltis applanata*, 130.
- Microsorium irregulare*, 561.
- Miers, J., on *Napoleona*, *Omphalocarpum*, and *Asteranthos*, 1; on the *Auxemmeæ*, a new Tribe of the *Cordiaceæ*, 23; on the *Barringtoniaceæ*, 47.
- Mitraria Commersoni*, 68.
- Mode of growth, of *Ballia callitricha*, 211; of *Corallinaceæ*, 203.
- Monachosorum davallioides*, 546.
- Monocotyledons*, self-fertilization sp. of, 378.
- Montbretia abyssinica*, 272.
- Moraea andongensis*, 271; *candelabrum*, 271; *glutinosa*, 271; *gracilis*, 272; *spithamæa*, 271; *textilis*, 270; *Welwitschii*, 270.
- Morphology of Primulaceæ, Dr. M. T. Masters on, 285.
- Mortierella*, fructification of, 421.
- Mucedines* (of Brisbane), 406.
- Mycelia* (of Brisbane), 407.
- Mycenastrum corium*, 406.
- Mycoides parasitica*, Dr. D. Douglas Cunningham on, 301; alliances of, 311; descrip. of, 301; disk and filaments, 304, 305, 310; gen. charac., 312; life-history of, 301; oogonium and zoospores of, 308; relations when attacked by lichens and fungi, 312; sexual fructification of, 307; spores of, 306; zoospores of, 309.
- Myosotis*, convolute corolla of, 177; scorpioid racemes of, 616.
- Myrtus alata*, 93.
- Myxogastres*, 128; of Brisbane, 406.
- Napoleona*, analysis of, 2; calyx of, 3; corona of, 2, 5; fruit of, 6; ovary of, 4; seeds of, 4.
- *angolensis*, 6, 12; *cuspidata*, 10; *Heudelotii*, 4, 8, 21; *imperialis*, 6, 7, var. *purpurea*, 8; *Mannii*, 11; *Vogelii*, 6, 9; *Whitfieldii*, 6, 9.
- Napoleonæ*, position and history of, 1.
- Nectria Eugeniæ*, 130.
- Nephrodium abruptum*, 532, 534; affine, 520; *amboinense*, 532; *angustifrons*, 527; *apiciflorum*, 517, 602, var. *nidus*, 518; *appendiculatum*, 516; *arbuscula*, 532; *aridum*, 531, 601, 602; *artinexum*, 536; *barbigorum*, 522; *Beddomei*, 517; *Boryanum*, 527, 528, 600, 603, var. *microstegioides*, 528, 544; *Brunonianum*, 522, 602; *Buchanani*, 525; *calcarata*, 515; *canum*, 515, 543, 602, 603; *cicutarium*, 539, 540, 603, var. *coadunata*, 540; *ciliatum*, 514, 602, 603; *Clarkei*, 518, 520, 536; *coadunatum*, 540; *cochleatum*, 520, 521, 603; *crenatum*, 524, 602; *crinipes*, 533, 534; *cucullatum*, 530, 531, 601, 602; *cuspidatum*, 512, 601; *da-*

- currens*, 539; *delicatulum*, 540; *didymosorum*, 530; *dissectum*, 526, 527, 534, 535; *divisum*, 527; *elatum*, 531, var. *procerum*, 531; *elongatum*, (*Swtz.*) 513, (*Hk. & Grev.*) 520; *Elwesii*, 516; *eriocarpum*, 525; *cursorum*, 534; *exaltatum*, 541; *extensum*, 516, 529, 530, 533, var. *laete-repens*, 530, var. *microsora*, 530, var.  $\beta$ . *minor*, (*Bedd.*) 532; *Fairbankii*, 517; *falcilobium*, 515, 517, 602, 605, var. *pubera*, 515; *Falconeri*, 522; *ferox*, 534; *fibrillosa*, 518; *Filix-Mas*, 519, 527, var. (*Bedd.*), 518, var. *cristata*, 519, (*Rich.*) var. 602, var. *fibrillosa*, 520, var. *intermedia* (*Bedd.*), 522, var. *khasiana*, 519, var. *marginata*, 520, 521, var. *normalis*, 519, var. *panda*, 519, var. *patentissima*, 520, var. *Schimperiana*, 520, (*Hochst.*) 522; *flaccidum*, 513, 514, 528; *fuscipes*, 526, 527, 535, 538; *gigantum*, 539; *glandulosum*, 531, 547, var. *laete-strigosa*, 532; *graciliscens*, 513, 517, 528, var. *decipiens*, 514, var. *hirsutipes*, 514, var. *didymochlaenoides*, 514; *grandifolium*, 537; *heterosorum*, 537, 555; *hirsutum*, 524; *hirtipes*, 513, (*Hk.*) var., 603; *Hudsonianum*, 534; *immersum*, 514; *ingens*, 526, 527, 535; *lacerum*, 522; *latipinna*, 532; *Leuzcanum*, 535, 536, 605; *lineatum*, 547; *lingua*, 553; *maderense*, 521; *membranifolium*, 534, 602, 603, var. *typica*, 535, var. *dimorphum*, 535; *meniscioides*, 532; *microphyllum*, 584; *microstegium*, 544; *multicaudatum*, 427, 540, 603; *multijugum*, 533, 602; *nymphale*, 533; *oethodes* (*Hk.*), 516; *odontoloma*, 521, var. (*Baker*), 518; *odoratum*, 525; *pallidum*, 523; *parasiticum*, 516, 532, 533, 534, 601, 602, 605, var. *aurea*, 533, var. *multijuga*, 533; *Parishii*, 525; *patentissimum*, 520; *pennigerum*, 532, var. *multilineata*, 532; *platypus*, 528; *polymorphum*, 537, 603, var. *Simonsii*, 537; *procurrens*, 530; *prolixum*, 516, 517, 602; *propinquum*, 529; *pteroides* (*J. Sm.*), 529; *pteridioides* (*Griff.*), 518; *puberulum*, 513, 514; *pulvinuliferum*, 525, 526, 602; *purpurascens*, 523; *remotum* (*Hk.*), 521, 523, var. *Chanteriae*, 523; *repandum*, 537; *rhodolepis*, 427, 526; *rigidum*, 523; *rostratum*, 538; *sagenioides*, 535, 602; *scabrosum*, 527, 602; *setigerum*, 528, 546; *siifolium*, 538; *sikkimense*, 525; *sparsum*, 489, 519, 523, 605, (*Don*) var., 603, var. *gracilis*, 508, var. *latisquama*, 524, var. *membranaceum*, 489, var. *minor*, 524, var. *nitidula*, 524, var. *squamulosa*, 524; *spectabile*, (*C. B. C.*) 526, (*Hk.*) 518, 603; *spinulosum*, 523; *splendens*, 527; *subconfluens*, 536; *syrmaticum*, 518; *tenericaule*, 528, 545, 602, 605; *terminans*, 529; *Thelypteris*, 514, 517, 544; *truncatum*, 534, 602; *tuberosa*, 540; *undulatum*, 524; *unitum*, 529, (*R. Br.*) 601, (*Sieber*) 530; *variolosum*, 538, 603; *vastum*, 536, 603; *venulosum*, 534; *Wightii*, 427, 538.
- Nephrolepis acuta*, 541; *cordifolia*, 540, 603, 605; *exaltata*, 541, 602, 603; *volubilis*, 541, 603.
- Neuronia asplenioides*, 542.
- New British Lichens, by Rev. W. A. Leighton, 145, 237.
- New Zealand, contribution to the Lichenographia of, 275.
- Nidulariacoi*, 128; of Brisbane, 405.
- Niphobolus* (subgen.), 552; *adnascens*, 552, 598; *angustatus*, 559; *caudatus*, 552; *costatus*, 553; *detergibilis*, 555; *elongatus*, 552; *fissus* (*Bl.*), 554; *floccigerus*, 554; *laevis*, 552; *macrocarpus*, 559; *nummulariaefolius*, (*Bedd.*) 576, (*J. Sm.*) 554; *porosus*, 554, *Schmidianus*, 554; *sphaerocephalus*, 559; *sticticus*, 554; *subfurfuraceus*, 553; *varius*, 552; *venosus*, 553.
- Notholaena carnosae*, 575, 598; *Marrantae*, 567; *persica*, 455; *piloseloides*, 575, 598; *undulata*, 598.
- Nullipores* closely allied to *Corallines*, (footnote) 207.
- Nymphaceae*, self-fertilization sp. of, 349.
- Odontoloma Boryanum*, 451; *repens*, 451.
- Odontotrema longius*, 239; *majus*, 239; *minus*, 239.
- Cecidiacei*, of Brisbane, 407.
- Cecidium apocynatum*, 407.
- Oidium lactis*, filaments of, 421.
- Oleandra Cumingii*, 542; *musaeifolia*, 542; *neriiformis*, 541; *Wallichii*, 542, 603.
- Olfersia triquetra*, 473.
- Omphalocarpum*, 1; description and position of, 13; *elatum*, 16; *procerum*, 15.
- Onagraceae*, plumes on seed of, 165; self-fertilization sp. of, 363, 392.
- On *Adoxa Moschatellina*, by Rev. G. Henslow, 194.
- On a Collection of Fungi made by Mr. Sulpiz Kurz, by F. Currey, 119.
- On a new Species of *Helvella*, by W. Phillips, 423.
- On *Mycoidea parasitica*, a new genus of parasitic Algæ, and the part which it plays in the formation of certain Lichens, by Dr. D. D. Cunningham, 301.
- On *Napoleona*, *Omphalocarpum*, and *Asteranthos*, by J. Miers, 1.
- On origin of tertiary and quinary symmetry of Flowers with indefinite and spirally arranged members, by Rev. G. Henslow, 194.
- On some points in the Histology of certain species of *Corallinaceae*, by Major-Gen. R. J. Nelson and Prof. Duncan, 197.
- On some points in the Morphology of the *Primulaceae*, by Dr. Maxwell T. Masters, 285.
- On the African species of the genus *Coffea*, Linn., by W. P. Hiern, 169.
- On the Geographical Distribution of the *Meliaceae*, by M. Casimir de Candolle, 233.
- On the Growth of the Flower-stalk of the Hyacinth, by A. W. Bennett, 139.
- On the Hygroscopic Mechanism by which certain Seeds are enabled to bury themselves in the ground, by F. Darwin, 149.
- On the minute Structure and mode

- of Growth of *Ballia callitricha*, by W. Archer, 211.
- On the Nature of the Corolla of *Primula*, by Rev. G. Henslow, 195.
- On the occurrence of Conidial Fructification in the *Mucorini*, illustrated by *Choanephora*, by Dr. D. D. Cunningham, 409.
- On the origin of Floral Æstivations, with Notes on the Structure of the Cruciferous flower, on that of *Adoxa*, and on the Corolla of *Primula*, by the Rev. G. Henslow, 177.
- On the Origin of the prevailing systems of Phyllotaxis, by the Rev. G. Henslow, 37.
- On the Origin of the so-called Scorpioid Cyme, by the Rev. G. Henslow, 613.
- On the Self-fertilization of Plants, by the Rev. G. Henslow, 317.
- On the Symmetry of a Cruciferous flower, by Rev. G. Henslow, 191.
- Onoclea orientalis*, 434.
- Onychium auratum*, 458, 596; *capense*, 459; *japonicum*, 459, 596, var. *intermedia*, 459, var. (*Kurz*) 458, 459, var. *multisecta*, 459; *lucidum*, 459; *multisectum*, 459, 597.
- Opegraphia saxicola*, 250.
- Ophioglossum Aitchisoni*, 586; *circinatum*, 583; *cordifolium*, 586, 596; *filiforme*, 584; *flexuosum*, 583, 584; *japonicum*, 585; *laciniatum*, 587; *pedatum*, 583; *pendulum*, 586, 587; *polyphyllum*, 586; *reticulatum*, 586, 596; *scandens*, 584; *vulgatum*, 586, var. *Aitchisoni*, 586.
- Orchidaceæ, self-fertilization sp. of, 378.
- Ornithogalum benguelense*, 248; *capifolium*, 248.
- Orobanchaceæ, self-fertilization sp. of, 373.
- Osmunda Claytoniana*, 582, 596; *crispa*, 459; *interrupta*, 582, 583; *japonica*, 583; *lanceolata*, 579; *Leschenaultiana*, 583, 596; *Lunaria*, 587; *monticola*, 582, 596; *regalis*, 583, 596; *speciosa*, 583, 596; *virginiana*, 582; *zeylanica*, 587.
- Owonia*, 235.
- Palicourea (Aubl.)*, 169.
- Papaver*, 178, 180; *orientale*, 180.
- Papaveraceæ, self-fertilization sp. of, 349.
- Paradigma, history and generic character of, 30, 31; *Galeottiana*, 31.
- Parasitic alga (*Mycoidea parasitica*), Dr. Cunningham on, 301.
- Parkeria pteridioides*, 471.
- Passifloraceæ, self-fertilization sp. of, 366.
- Patagonula*, 28; history of, 27. — *americana*, 28; *bahiensis*, 30; *glabra*, 29; *Tweediana*, 29.
- Patania Elwesii*, 436.
- Pavetta (Linn.)*, 169.
- Pelargonium*, growth of flower-stalk of, 143.
- Pellæa calomolanos*, 461; *geraniæ-folia*, 597; *gracilis*, 460; *nitidula*, 455, 460; *nudiuscula*, 455; *Tamburii*, 460.
- Peranema aspidioides*, 434; *cyatbeoides*, 435, 599.
- Perigara globosa*, 86; *valida*, 94.
- Pertusaria graphica*, 275; *incarnata*, 241.
- Petersia*, 107; *africana*, 108; fruit of, 48.
- Peziza aurantia*, 128; *cinereonigra*, 404; *rutilans*, 128; *scutellata*, 404; *vinoso-brunnea*, 404.
- Phalloidei, of Brisbane, 405.
- Phegopteris (subgen.)*, 542; *Dryopteris*, 545; *luxurians*, 548; *opaca*, 568; *polypodioides*, 544; *Robertiana*, 545; *rugulosa*, 546; *stegogramme*, 569; *Totta*, 567; *urophylla*, 547; *vulgaris*, 544.
- Phillips, W., now species of *Helvella*, 423.
- Phlyctis, neo-zelandiæ*, 281; *ocellata*, 281; *sordida*, 281.
- Phorolobus crispus*, 459.
- Phycopeltis*, 315.
- Phylloidy of *Primulaceæ*, 286.
- Phyllotaxis, Airy's condensation-theory of leaf-arrangement criticized, (footnote) 38; correlation between internodes and angular divergences, 40; di- and tristichous and pseudo-distichous arrangement, 43; divergence  $\frac{8}{27}$  determines primary series, 41; law of angular distance, 38; of *Lepidodendron* (fossil), 40; opposite leaves fail to produce  $\frac{1}{2}$  and  $\frac{1}{3}$  arrangement, 42; origin of cycles, 39; origin of prevailing system of, Rev. G. Henslow on, 37; reduction from whorled leaves, (footnote) 44; relative ages, di- and monocotyledons, 44; results of arrest of development, 44; scheme of ordinary phyllotaxis, 44; spiral arrangement deduced from opposite and decussate leaves, 37; theoretical conditions, 41.
- Phymatodes (subgen.)*, 557; *cyrtolum*, 564; *polycarpa*, 561.
- Pinguiculaceæ self-fertilization sp. of, 377.
- Piptocephalis*, fructification of, 420.
- Pistil, as adapted to self-fertilization, 380.
- Placodium argillaceum*, 282; *illitum*, 282; *lecanorinum*, 282; *Rhoderi*, 282.
- Plagiogyria adnata*, 472; *glauca*, 472; *pyncophylla*, 473; *scandens*, 472; *triquetra*, 473.
- Planchonia, charac. and history of*, 90. — *crenata*, 91; *elliptica*, 93; *littoralis*, 94; *nitida*, (*Bl.*) 93, (*Miq.*) 88; *sundiaca*, 81, 92; *tetraptera*, 93; *undulata*, 92; *valida*, 81, 94.
- Plants, self-fertilization of, by the Rev. G. Henslow, 317.
- Pleocnemia (subgen.)*, 534; *Clarkei*, 536; *javanica*, 535; *Luzeana*, 535.
- Pleopeltis (subgen.)*, 566; *capitellata*, 566; *dilatata*, 566; *ebenipes*, 564; *Griffithiana*, 560; *hemionitidea*, 562; *heterocarpa*, 561; *himalayensis*, 566; *incana*, 596; *irioides*, 561; *Lehmanni*, 566; *leiorhizon*, 567; *longissima*, 565; *malacodon*, 563, 564; *normalis*, 558; *nuda*, 558; *ovata*, 560; *oxyloba*, 563; *Parishii*, 564; *punctata*, 561; *rhyncophylla*, 560; *Stewartii*, 563; *superficialis*, 557; *tridactyla*, 562.
- Plethostephia*, 32; *angiocarpa*, 32.
- Pœcilopteris costata*, 580; *flagellifera*, 580; *terminans*, 581.
- Polybotrya appendiculata*, 577; *intermedia*, 577; *marginata*, 577;

- neglecta, 578; nodiflora, 577; virens, 581.
- Polygalæ, self-fertilization sp. of, 352.
- Polygonacæ, self-fertilization sp. of, 378, 393.
- Polypodium acrostichoides, 513, 552; aculeatum, 509; acutissimum, 558, 605; adhærens, 605; adnascens, 552, 600, 601; adnatum, 602; alternans, 602; alternifolium, 565, 601; altissimum, 433; amœnum, 550, 601, var.? tonglensis, 550; angustatum, 559, 601, var. depauperata, 559; appendiculatum, (Bedd.) 543, (Bk.) 516, var. squamæstipes, 543; arboreum, 605; argutum, 551, 552, 601; aristatum, 511; atropunctatum, 558; auriculatum, 433, 543, (L.) 507, (Wall.) 569, 601; Barometz, 435; Blumei, 570; Boothii, 555; brauneum, 544, 602; calcareum, 545; capitellatum, 566, 601; caudatum, 552; caudigerum, 530, 601; chatagrammicum, 427, 548, 549; chinense, 558; cicutarium, 539; clathratum, 427, 559; comosum, 602; confluens, 602; conifolium, 546, 601; conjugatum, (Hk. & Bk.) 557, (Lamk.) 557; contaminans, 432, 602; contiguum, 557, 558, 601; cordifolium, 540; coriaceum, 559; coronans, 556, 557, 601; costatum, 553, 555, 600; crenatum, 525; curvinervium, 601; cyrtolobum, 563, 564; darexforme, 427, 443; davallioides, 546; dentigerum, 491, 602; detergibile, 555; dichotomum, 428; dilatatum, 536, 565, 601; distans, 544, 568, 601, var. adnata, 544, var. glabrata, 544, var. minor, 545; Dodgsoni, 563; drepanopterum, 493, 494; Dryopteris, 545; ebenipes, 563, 564, 601, var. Oakesii, 564, var. Parishii, 564; ellipticum, 570; elongatum, 513, 601; erubescens, 516, 543, 598, 602; erythrocarpum, 427, 565; exaltatum, 541; excavatum, 558, 559, 565; falcatum, 512; Fieldingianum, 551; Filix-Mas, 519; Finlaysonianum, 605; fissum, 554, 600, var. floccigerum, 554; flocculosum, 554, 555, 600; furfuraceum, 601; fragrans, 454; giganteum, 433, 601; glabrum, 561, 601; gladiatum, 558, 601; glaucistipes, 601; glaucum, 428; grandifolium, 560, 601; Grevillecanum, 558, 605; Griffithianum, 560; Griffithii, 544; gibbosum, 549; hastatum, 562, 564, 601, var. oxyloba, 563, var. Thunbergii, 563; hemionitideum, 561, 601; Hendersoni, 550; heteractis, 553, 554; heterocarpum, 560, (Bl.) 561, 562, (Hk. & Bk.) 561; himalayense, 566; Hippocrepis, 539; Horsfieldii, 601; hymenodes, 557, 579, 601; hyperboreum, 434; involutum, 570; irioides, 603; jaintense, 427, 552; javanica, 535; juglandifolium, 566, 601, var. biserialis, 566; khasyanum, 549; Kulhaitense, 493, 531; lachnopus, 550, 551, 601; lanceolatum, 560; latebrosum, 432; Lehmanni, 566; leiopteris, 558; leiorhizon, 566, 567, 601; Leschenaultianum, 602; Leuzeanum, 535; Linnæi, 556, 601; Lindleyanum, 601; lineare, (Burm.) 428, 558, 559, (Thunb.) 601, 605, var. polymorpha, 559, var. stenisto, 559; lineatum, 548, 601, var. penangiana, 548; lingua, 555, var. (Hk. & Bk.), 553; longifolium, (Bl.) 601; longifrons, 558, 601; longipes, 544, 601; longissimum, 556, (Mett.) 565; lonchites, 506; loriforme, 558, 559, 601; loxogramme, 570; lucidum, 567; luxurians, 548; macrodon, 536; malacodon, 564; marginale, 447, 602, (Wall.) 546, 527, 601; mauritianum, 601; melanopus, 564; membranaceum, 560, 601; meniscioides, 547; microrrhizoma, 551, 601; molle, 533; molliusculum, 533, 602; mouleinsis, 566; multilineatum, 494, 547, 548; mysurense, 554, 600; nemorale, 533, 601; neriifolium, 601; nigrescens, 565, 601, var. polyphylla, 601; normale, 557, 558, 601; nummulariaefolium, 553, var. obovatum, 554; obliquatum, 549; oppositum, 524, 605; ornatum, 528, 545, 546, 601; ovatum, 560, 601; oxylobum, 601; oxyphyllum, (Wall.) 486, 493, 602, (Kze.) 561; pallidum, 545, 546; paludosum, 544; palustre, 577; parasiticum, 533; Parishii, 564; parvulum, (Bedd.) 549, (Bory) 549; pedunculatum, 570; pertusum, 552, 600; phegopteris, 544; phlebodes, 558, 559; phymatodes, 557, 602; polycephalum, 561, 601; polyodon, 512, 605; polyphylla, 565; polypodioides, 544; porosum, 554, 600; pothifolium, 571; proliferum, 548, 601; propinquum, 556, 563, 600, var.? (Wall.), 563; pteropus, 562, 601; pubigerum, 535, 605; punctatum, (Swetz.) 601, (Thunb.) 528, 546, 561; quercifolium, 555, 556, 557, 601; rhyncophyllum, 560; rivale, 556; Robertianum, 545; rostratum, 557; rubrinerve, 547; rugulosum, 545, var. hirsutum, 546; Russellianum, 528, 605; scabridum, 531; scabrum, 605; Scottii, 542; secundum, 529, 601; semibipinnatum, 605; serra, 601; serriola, 549; sesquipedale, 558, 601; siifolium, 538; sinuosum, 605; sophoroides, 605; Speluncæ, 448; sphaerocephalum, 559, 601; stenophyllum, 559; Stewartii, (C. B. C.) 563, (Bedd.) 563; stipitum, 544; stigmatosum, 553, 600; subamœnum, 427, 550; subarborescens, 605; subauriculatum, 551; subdigitatum, 546; subfalcatum, 549; subfurfuraceum, 553, 555; subpectinatum, 601; subtripinnatum, 427, 545; superficiale, 557, var. semilinearis, 558; tenericaule, 546, 602, (Hk.) 566, (Wall.) 528; tenerifrons, 549; Thelypteris, 544; tomentosum, 555; Totum, 567; trichodes, 546; trichomanoides, 549, 550; tridactylon, 562, 601; trifidum, 563; triplum, 563; truncicola, 549; umbrosum, 433, 632; unitum, 529; urophyllum, 547, 571, 601, var. khasiana, 547; varium, 552; venosum, 605; venustum, (Wall.) 566, (Desv.) 567, 601, var. niphoboloides, 567; verrucosum, 552,

- 601; viscosum, (*Roxb.*) 546, (*Zipp.*) 546; vittarioides, 552, 601; Wallichii, 555, 601; Wightianum, 558, 605; Willdenovii, 556; Zippellii, 561, (*Bl.*) 562, (*Mett.*) 562; zosteræforme, 601; Zollingerianum, 562.
- Polyporei, 121; of Brisbane, 400.
- Polyporus affinis, 122, 401; amboinensis, 122; anebus, 124; applanatus, 123; Broomei, 402; brunneus, 400; brunneo-pictus, 122; calcens, 402; eichoraceus, 402; cinerascens, 124; cinereo-fuscus, 124, 402; cinnabarinus, 124; crassipes, 122; Feoi, 124; ferreus, 124; flabelliformis, 122, 401; floridus, 121; fruticum, 401; funalis, 123, 402; fuscolineatus, 401; hirsutus, 403; holosclerus, 123; hypoplastus, 121; ignarius, 401; incertus, 123; lucidus, 122, 401; luteo-nitidus, 400; luteo-olivaceus, 402; marginatus, 123; modestus, 122; nilghericensis, 122; perennis, 121; Persoonii, 123; picipes, 122, 401; pinsitus, 124; platotus, 401; quadrans, 400; rhipidium, 401; rubidus, 123, 402; sanguineus, 122, 401; seruposus, 124; senex, 401; Splitgerberi, 122; versicolor, 124; xanthopus, 121, 400; xerophyllaceus, 124.
- Polysaccum olivaceum, 406; pisocarpium, 406.
- Polystichum (subgen.), 585; aculeatum, 509; aristatum, 511; Atkinsoni, 506; auriculatum, 489; ilicifolium, 506, var. minimum, 506, var. minus, 506; Lachenense, 506; Lonchitis, 506; Prescottianum, 510; radicans, 507; sikkimense, 525; stimulans, 508; Thomsoni, 508, 509; truncatum, 534.
- Porina endochrysa, 280.
- Portulacæ, self-fertilization sp. of, 392.
- Preliminary Note on Growth of Female flower-stalk of *Valisneria spiralis*, by A. W. Bennett, 133.
- Primroses, double, 287, 293.
- Primula, corolla of, Rev. G. Henslow on, 177.
- , æstivation and diagrams of, 177, 189; nature of the corolla of, 195; cupped petals of, 289.
- Primula cortusoides, 286; japonica, 286; prænitens, 286, sinensis, 286; vulgaris, 189.
- Primulacæ, androecium of, 289; bibliography of, 297; calyx of, 285; chorisis of, 288; corolla of, 286; develop. parts of flower, 287; enation of, 288, 289; gynæcium of, 292; morphology of, Dr. Masters on, 285; ovules of, 295; pistillody, stamens of, 290; placentation of, 292, Trécul's view of, (ftnote) 293; self-fertilization sp. of, 376, 393; some gen. and sp. subject to changes, 727; summary of morphology, 296; tubular petals in, 289; variations in calyx of, 286.
- Prismatomeris (*Thw.*), 169.
- Prunus, calycine lobes of, 196.
- Psathyra, 119.
- Psathyrella, 119.
- Pseud-Allantodia (subgen.), 495.
- Psilosiphon cyanescens, 272; fragrans, 272; odoratissimus, 273.
- Psilotum flaccidum, 596; nudum, 589, 596; triquetrum, 589, 596.
- Psychotria (*Linn.*), 169.
- Pteris acuminatissima, 461; alternifolia, 604; amplexicaulis, 461, 597; anamallayensis, 469; angustata, 597; angustifolia, 574; aquilina, 468, 597, 604, var. esculenta, 468, var. lanuginosa, 468; argentea, 458; arguta, 604; aspericaulis, 465, 467, 470, 597; aurita, 471; australis, 595; brevisora, 468; biaurita, 469, 470, 597, var., 604, var. geminata, 469; bicolor, 457; Blumeana, 466; brevisora, 467, 468; cæspitosa, 455, 597; calome-lanos, 461; caudata, 468; chrysocarpa, 458; costata, 461; crenata, 462, 463, 467; croatica, 452, 462, 463, 464, 597; crispa, 459; dactylina, 463; Dalhousiæ, 465; densa, 468, 597; digitata, 463, 597; dimidiata, 464; diversifolia, 461; ensiformis, (*Burm.*) 463, 464, 604, 605, (*Ham.*) 605, var. Grevilleana, 464; esculenta, 468; excelsa, 467, 597; farinosa, 457; firma, 468, 597; flabellata, 464; geminata, 604; gracilis, 455, 460; graminifolia, 574; Grevilleana, 464, 466, 605; Griffithii, 464; hastata, 461, 597; heterodactyla, 464; incisa, 471, 597; læta, 462, 597; lanigera, 468; lanuginosa, 468, 597; linearis, 597; longifolia, 461, 597; longipes, 465, 468; longipinnula, 466, 467, 469, 597; lorigera, 467, 468, 597; ludens, 470, 597; lunulata, 452; marginata, 471; multidentata, 463, 605; multifida, 470; mysurensis, 470, 597; nemoralis, (*Hk. & Bauer*) 465, (*Willd.*) 469, 597; nervosa, 462, 597; nitidula, 460, 597; palmata, 470; pectinata, 465, 470; pedata, 470; pellucens, 468, 470; pellucida, 462, var. stenophylla, 463, var., 597; piloselloides, 575; polita, 597; pyrophylla, 465; quadriaurita, 465, 466, 467, (*Hk.*) 467, (*Retz.*) 467, 469, 470, 597, var. Blumeana, 466, var. khasiana, 466, var. major, 466; radiata, 505; recurvata, 468, 597; repandula, 466; revoluta, 468; revolvens, 471; sagittifolia, 471; scabra, 605; scabripes, 463, 597; scandens, 577; semihastata, 467, 468, 597; semipinnata, 464, 597; setigera (or var.), 465; sinuata, 471, 597; spinescens, 465; Stelleri, 460; subquinata, 465, 467; succulenta, 471; subindivisa, 427, 467; subpedata, 597; terminalis, 467, 597; tripartita, 470; umbrosa, 470, 597; varians, 455, 597; venulosa, 462; vespertilionis, 471; vittata, (*L.*) 461, (*Roxb.*) 461; Wallichiana, 468, 469, 470, 597, var. quadri-pinnata, 470; Wightiana, 468, 604; Zollingeri, 468.
- Queensland, fungi of, 399.
- Randia genipæflora (*DC.*), 169.
- Ranunculacæ, self-fertilization sp. of, 347, 391; twisting seed and awn of, 164.

- Ranunculus bulbosus*, aestivation of, 189.
- Report on the Liliaceæ, Iridaceæ, Hypoxidaceæ, and Hamodoraceæ of Welwitsch's Angolan Herbarium, by J. G. Baker, 245.
- Review of the Ferns of Northern India, by C. B. Clarke, 425.
- Rhizophora*, embryo of, 52.
- Rhizomorpha Harrimanni*, 407.
- Rhizophoraceæ, remarks on, 54.
- Rhododendron Griffithianum*, 19; maximum, 19.
- Ribes coccineum*, aestivation of, 190.
- Romneyceæ, 180.
- Rosaceæ, self-fertilization sp. of, 362, 392.
- Rubiaceæ, self-fertilization sp. of, 366, 392.
- Rudgea (Salisb.)*, 169.
- Sacellium*, gen. charac. of, 25; lanceolatum, 25.
- Sagenia alata*, 536; *cicutaria*, 535; *coadunata*, 540; *gigantea*, 539; *heterocarpa*, 537; *hippocrepis*, 539; *macrodonata*, 539; *pteropus*, 539; *silhetensis*, 540; *variolosa*, 540.
- Salvinia natans*, 598.
- Samolus Valerandi*, 286; variation corolla of, 287.
- Sandersonia littonioides*, 262.
- Sandoricum indicum*, 236; (*Main-gayia*) *marginatum*, 236.
- Sansevieria bracteata*, 253.
- Saxifragaceæ, self-fertilization sp. of, 365.
- Schizaea dichotoma*, 595; *digitata*, 583, 595.
- Schizobasis angolensis*, 255.
- Schizoloma ensifolium*, 452; *Griffithianum*, 452.
- Schizophyllum commune*, 121, 400.
- Scilla arenaria*, 249; *benguellensis*, 249; *congesta*, 250; *flaccidula*, 249; *hispidula*, 248; *lanceifolia*, 250; *laxiflora*, 250; *platyphylla*, 250; *polyphylla*, 249; *simiarum*, 249.
- Sclerodesma Bovista*, 406.
- Scelopendrium Ceterach*, 504; *dubium*, 496.
- Scorpioid cyme, Rev. G. Henslow on, 613.
- Scorpioid cyme as exemplified in *Lagerstroemia*, 613; critical remarks on, 617; description of, 613; Henslow's views, 613; in *Silene pendula*, 613; in *Lathræa Squamaria*, 613; Kaufmann and Warming's views, 620; oscillating arrangement, 613; Payer's definition, 619; Sachs's view, 619; summary of successive steps to, 618.
- Scrophulariaceæ, self-fertilization sp. of, 370, 393.
- Seeds burying themselves, hypothesis on, 166; hygroscopic mechanism of, 149.
- Self-fertilization of Plants, Rev. G. Henslow on, 317.
- , advantages derived from, 319; are fittest to survive struggle for life, 396; are perfectly healthy, 388; chief facts tabulated, 322; cleistogamous flowers self-fert., 347; conservation of energy in reduct. stamens and pollen, 380; Darwin's views on, 320; fertility and health does not decrease, 382; fertility may increase, 383; few flowers (physiolog.) self-sterile, 323; flowers may become self-fertilizing in absence of insects for intercrossing, 331; free from competition self-fertiliz. may equal intercrossed, 383; grafting effects of, 333; highly self-fert. var. arise under cultivation, 339; inconspicuous flowers self-fert. or else anemophilous, 342; introductory remarks on, 317; loss of colour correlated with self-fertilization, 327; origin of conspicuous flowers and causes inducing reversion to self-fertilization, 333; majority of flowers self-fertile, 323; many plants morphol. self-sterile, 324; Mehan's views on, 320; more productive than flowers dependent on insects, 390; naturalized abroad, replace native vegetation, 395; partial and total arrest devel. of corolla, 327; partial excision of corolla and stamens, 326; physiol. and morphol. self-sterile plants may become self-fertile, 325; plants may become self-fertilizing on reduction of temperature, 332; relative fertility may surpass crossed plants, 382; secured by flowers closing after expansion, 329; some derive no benefit from crossing with same on distinct stock, 385; some flower-buds never open, 330; special adap. for self-fert. of Brit. plants, 347; summary of adaptations to secure self-fert., 380; value of intercrossing, 318; withering of corolla, 325; world-wide distrib. of self-fert. Brit. plants, 391.
- Selliguea decurrens*, 571; *Hamiltoni*, 570; *lanccolata*, 570; *homionitidea*, 562; *Wallichiana*, 570.
- Shuta*, native name of *Agasta splendens*, 60.
- Silene pendula*, scorpioid cyme of, 614.
- Sitobium strigosum*, 436.
- Smilacæ (Angolan), 264.
- Smilax Kraussiana*, 264.
- Solanaceæ, self-fertilization sp. of, 370, 392.
- Solanææ, scorpioidal infloresc. of, 616.
- Sphaclaria callitricha*, struct. of, 220.
- Sphæria phaselina*, 130; *sublimbata*, 130.
- Sphæriacei, 129; of Brisbane, 405.
- Sphæropteris barbata*, 435, 526, 599; *Hookeriana*, 434, 603.
- Sporocybacei, 129.
- Stamens as adapted to self-fertilization, 380.
- Stegnogramme aspidioides*, 569.
- Stemonitis fusca*, 406.
- Stenochlæna pycnophylla*, 473; *scandens*, 577; *triquetra*, 473.
- Stereum adustum*, 127; *cyathiforme*, 127; *elegans*, 126; *lobatum*, 126, 403; *medicum*, 127; *Ostrea*, 126; *papyrinum*, 127; *scytale*, 127.
- Sticta canaliculata*, 282.
- Stilbacei, of Brisbane, 406.
- Stilbum erythrocephalum*, 129; *inconspicuum*, 129.
- Stipa-awn*, 155, hygroscope, 156.
- Stipa pennata*, descrip. of seed, 149; hygroscopic action of awn of, 150; mechanism of torsion in awn of, 157.

- Stravadium*, 79, 80; *acuminatum*, 102; *acutangulum*, 80; *album*, 65; *angustatum*, 105; *excelsum*, 112; *coccineum*, 83; *cochinchi-nense*, 101; *costatum*, 88; *demis-sum*, 81, 83; *denticulatum*, 88; *globosum*, 86; *gracile*, 86; *Horsfieldii*, 85; *insigne*, 75; *in-tegrifolium*, 89; *lucidum*, 88; *lu-zonense*, 84; *macrophyllum*, 61; *obtusangulum*, 77, 81; *pubescens*, 83; *Reinwardtii*, 188; *reticulatum*, 87; *Rhecdii*, 82; *rubrum*, 68, 69; *Sarcostachys*, 102; *semisutum*, 89; *serratum*, 87; *spicatum*, 85.  
 Structure of *Ballia callitricha*, 211.  
*Struthiopteris orientalis*, 434.  
 Suppression of 5th part cycle of flowers, 192.  
*Swietenia Mahagoni*, 236.  
*Symplocineæ*, 1.  
*Syngamma vestita*, 568.  
  
*Tæniopsis lineata*, 574.  
*Tænitis blechnoides*, 575; *interrup-tus*, 575; *piloselloides*, 575; *pte-roides*, 575; *pusilla*, 574.  
*Thamnopteris Grevillei*, 476; *nidus*, 475; *phyllitidis*, 475; *Simonsiana*, 476.  
*Thecacophora globuligera*, 407.  
*Thelenella Wellingtonii*, 280.  
*Thelephora palmata*, 126; *pusilla*, 126.  
*Thelotrema monosporum*, var. *pa-tulum*, 278; *saxatile*, 278.  
*Tigellum*, term defined, 53.  
 Torsion in *Avena*, 163; of cells, 161; of seeds, 164; on mechanism in *Stipa*-awn, 157; power of, 161.  
*Torulacei*, of Brisbane, 406.  
*Trametes eingulatus*, 125; *lobatus*, 124; *occidentalis*, 125; *perennis*, 402; *rigida*, 402; *umbrinus*, 124.  
*Tremella foliacea*, 127.  
*Tremellini*, 127, of Brisbane, 404.  
*Tremellodon gelatinosum*, 126.  
*Trichia pyriformis*, 128.  
*Trichilia*, 235; *havanensis*, 236; *pe-ruviana*, distrib. of, 233.  
  
*Trichobasis rubigo-vera*, 407.  
*Trichocoma paradoxum*, 128.  
*Trichodermacei*, 126, 128.  
*Trichogastres*, of Brisbane, 406.  
*Tricholoma*, 119.  
*Trichomanes alchemillæfolium*, 598; *anceps*, 598; *auriculatum*, 441; *Belangeri*, 441; *bipunctatum*, 440; 598; *campanulatum*, 604; *chi-nense*, 449; *corticola*, 440; *den-ticulatum*, 438; *digitatum*, 440; *dissectum*, 441; *Filicula*, 440; *gracile*, 440; *insigne*, 440; *java-nicum*, 442, 598; *Kunzeanum*, 441; *Kurzii*, 440; *limbatum*, 441; *longisetum*, 598; *muscoides*, 439, var. *sublimbatum*, 439; *nanum*, 440; *parvulum*, 440; *plicatum*, 440; *proliferum*, 441; *pyrami-dalis*, 598; *pyxidiferum*, 440, var. *limbatum*, 441; *radicans*, 441, 442, 598, var. *anceps*, 441; *rigi-dum*, 442, 598; *Schmidianum*, 440; *setigerum*, 442, 598; *strigo-sum*, 447; *sublimbatum*, 439, 440; *umbrosum*, 441, 598; *undulatum*, 598; *villosa*, 598.  
*Trypethelium astroidea*, 278; *Cu-mingii*, 278; *pyrenuloides*, 278.  
*Tsjeria Saamstravadi*, 82.  
*Tulbaghia æquinoctialis*, 246.  
*Typhula fuscipes*, 127.  
  
*Ugena microphylla*, 584; *polymor-pha*, 584; *semihastata*, 584.  
*Umbelliferae*, self-fertilization sp. of, 365, 392.  
*Urceolaria novæ-zelandiæ*, 275.  
*Urginea psilostachya*, 247; *chloran-tha*, 248; *comosa*, 247.  
*Ustilago axicola*, 407; *carbo*, 407.  
  
*Valerianaceæ*, self-fertilization sp. of, 367.  
*Vallisneria spiralis*, growth of female flower-stalk of, 133.  
*Vavæa Amicorum*, 235.  
*Vellosia capillaris* (*Welw.*), 264; *squar-rosa* (*Welw.*), 264; *stenophylla* (*Welw.*), 265; *velutina* (*Welw.*), 265.  
*Verbenaceæ*, self-fertilization sp. of, 393.  
*Verrucaria aspistea*, 280; *astata*, 280; *circumpressa*, 278; *dealbata*, 280; *epidermidis*, var. *gemellipara*, 279; *Eschweileri*, 280; *fumosaria*, 239; *gemellipara*, 278; *Larbalestierii*, 242; *micromma*, 279; *minutis-sima*, 270; *myriospora*, 145; *ne-ottizans*, 239; *nitida*, 280; *oc-culta*, 279; *pertenuis*, 239; *pru-ino-grisea*, 279; *pyrenastroides*, 278; *saxicola*, 280; *subbiformis*, 279; *succina*, 145; *Thwaitesii*, 278.  
*Viburnum Tinus*, æstivation of, 189.  
*Violaceæ*, self-fertilization sp. of, 351.  
*Vittaria divergens*, 604; *elongata*, 573, (*Swtz.*) 598, (*Wall.*) 574; *falcata*, var., 574; *flexuosa*, 574; *interrupta*, 451; *japonica*, 574; *lineata*, 574; *minor*, 574, var. *minima*, 574; *plantaginea*, 574; *rigida*, (*Kaulf.*) 573, (*Wall.*) 574, 578; *sikkimensis*, 574; *zosteræ-fovia*, 574.  
*Volvaria*, 119.  
*Vutu-dina* (a *Barringtonia* ?), 57.  
  
*Walleria angolensis*, 262.  
*Welwitsch's Angolan Herbarium*, re-port on *Liliaceæ* &c. of, 215.  
*Woodsia elongata*, 435; *hyperborea*, 434, 435; *lanosa*, 435; *mollis-sima*, 435, f. *subcordata* (*Milde*), 434.  
*Woodwardia radicans*, 475, 595.  
  
*Xerophyta capillaris*, 264; *squarrosa*, 264; *stenophylla*, 265; *velutina*, 265.  
*Xylaria digitata*, 129; *flagelliformis*, 129; *guyanensis*, 129; *hypoxylon*, 129; *Kurziana*, 129; *mutabilis*, 129; *pileiformis*, 405; *polymorpha*, 129, 405; *rhytidophlea*, 405; *tabacina*, 129.